

## Approximation of the Series $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{an^2+bn+c}$

where  $a, b, c \in \mathbb{R}$  with  $a \neq 0$

**Kumari Sreeja S. Nair and Dr. V. Madhukar Mallayya**

*<sup>1</sup>Assistant Professor, Department of Mathematics  
 Govt. Arts College, Thiruvananthapuram, Kerala, India.*

*<sup>2</sup>Former Professor and Head, Department of Mathematics  
 Mar Ivanios College, Thiruvananthapuram, Kerala, India.*

### Abstract

Here we give approximation of an alternating series using remainder term of the series. Here we introduce a new term called correction term. The correction term plays a vital role in series approximation.

**Keywords:** Correction function, error function, remainder term, alternating series, rational approximation, Dirichlet's series.

### INTRODUCTION

The illustrious mathematician Madhava of 14<sup>th</sup> century introduces correction function for the series for pi. The Madhava series is

$C = \frac{4d}{1} - \frac{4d}{3} + \frac{4d}{5} - \dots + (-1)^{n-1} \frac{4d}{2n-1} + (-1)^n \frac{4d(2n)/2}{(2n)^2+1}$ , where C is the circumference of a circle of diameter d.

Here the remainder term is  $(-1)^n 4d G_n$  where  $G_n = \frac{(2n)/2}{(2n)^2+1}$  is the correction term. The introduction of the correction term improves the value of C and gives a better approximation for it.

### RATIONAL APPROXIMATION OF ALTERNATING SERIES $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{an^2+bn+c}$

where  $a, b, c \in \mathbb{R}$  with  $a \neq 0$  and  $\sqrt{b^2 - 4ac} \neq 2a$ .

The alternating series  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{an^2+bn+c}$  satisfies the conditions of alternating series test and so it is convergent.

#### Theorem

The correction function for the alternating series  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{an^2+bn+c}$  where  $a, b, c \in \mathbb{R}$  with  $a \neq 0$  is  $G_n = \frac{1}{\{2an^2+(2b+2a)n+(2c+b+2a)\}}$

#### Proof

If  $G_n$  is the correction function after  $n$  terms of the series, then

$$\text{we have } G_n + G_{n+1} = \frac{1}{an^2+(2a+b)n+a+b+c}$$

$$\text{The error function is } E_n = G_n + G_{n+1} - \frac{1}{an^2+(2a+b)n+a+b+c}$$

Let  $G_n(r_1, r_2) = \frac{1}{\{2an^2+(4a+2b)n+(2a+2b+2c)\} - (r_1 n + r_2)}$  where  $r_1, r_2 \in \mathbb{R}$  and  $n$  is fixed.

Then error function  $|E_n(r_1, r_2)|$  is minimum for  $r_1 = 2a$ ,  $r_2 = b$

Hence for  $r_1 = 2a$ ,  $r_2 = b$ , both  $G_n$  and  $E_n$  are functions of a single variable  $n$ .

Thus the correction function for the series  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{an^2+bn+c}$  is

$$G_n = \frac{1}{\{2an^2+(2b+2a)n+(2c+b+2a)\}}$$

The corresponding error function is

$$|E_n| = \frac{|(b^2-4ac)-4a^2|}{\{2an^2+(2b+2a)n+(2c+b+2a)\}\{(2an^2+(2b+6a)n+(6a+3b+2c))\}\{(an^2+(2a+b)n+(a+b+c))\}}$$

Hence the proof.

**REMARK**

Clearly  $G_n$  is less than the absolute value of the  $(n+1)^{\text{th}}$  term.

**APPLICATION**

1. The series  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2} = {}^n(2)$

We have  ${}^n(2) = 0.8224670334$ , using a calculator.

The correction function for the series is  $G_n = \frac{1}{2n^2+2n+2}$

For  $n=10$ , the series approximation after applying correction function is given below

Number of terms	$S_n$	$S_n + (-1)^n G_n$
10	<b>0.8179621756</b>	<b>0.82246666801</b>

2. THE ALTERNATING SERIES  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n(n+1)}$

The alternating series  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n(n+1)}$  is convergent and converges to  $2\log 2 - 1$ .

We have  $2\log 2 - 1 = 0.3862943611$ , using a calculator.

The correction function for the series is  $G_n = \frac{1}{2(n+1)^2+1^2}$

For  $n=10$ , the series approximation after applying correction function is given below

Number of terms	$S_n$	$S_n + (-1)^n G_n$
10	<b>0.3821789321</b>	<b>0.3863283098</b>

**CONCLUSION**

The introduction of correction function improves the sum of the series and gives a better approximation.

**REFERENCES**

- [1] Dr. Konrad Knopp - Theory and Application of Infinite series - Blackie and son limited (London and Glasgow)
- [2] Sankara and Narayana, Lilavati of Bhaskaracharya with the Kriyakramakari, an elaborate exposition of the rationale with introduction and appendices (ed) K.VSarma (Visvesvaranand Vedic Research Institute, Hoshiarpur) 1975, p, 386-391.
- [3] Dr. V.Madhukar Mallayya- Proceedings of the Conference on Recent Trends in Mathematical Analysis- © 2003, Allied Publishers Pvt. Ltd. ISBN 81-7764-399-1
- [4] A Course of Pure Mathematics - G.H.Hardy (tenth edition) Cambridge at the university press 1963
- [5] K. Knopp, Infinite sequences .and series, Dover-1956
- [6] T.Hayashi, T.K.Kusuba and M.Yano, Centaurus,33,149,1990
- [7] Yuktidipika of Sankara (commentary on Tantrasangraha), ed. K.V.Sarma, Hoshiarpur 1977