

# New Classes of General Quasi Variational Inclusions

Netra Kumar Gupta<sup>\*1</sup>, Muhammad Aslam Noor<sup>†2\*</sup>, Khalida Inayat Noor<sup>‡3</sup> and Suja Varghese<sup>§4</sup>

<sup>1</sup>*Department of mathematics Govt. V.Y.T.P.G.Autonomous College, Durg 491001 (C.G.), India.*

<sup>2,3</sup>*Mathematics Department, COMSATS University Islamabad, Islamabad, Pakistan.*

<sup>4</sup>*Department of mathematics ST. Thomas College, Bhilai, Durg 490006 (C.G.), India.*

## Abstract

In this article, we introduce and consider some new classes of general quasi variational inclusions. Special cases are discussed as applications.

We prove that the general quasi variational inclusions are equivalent to the fixed point problems. This alternative formulation is used to discuss the existence of a solution as well as to propose some iterative methods. Convergence analysis is investigated only using the Lipschitz continuous of the operators. Since the general quasi variational inclusions include quasi variational inequalities, variational inequalities, and related optimization problems as special cases, our results continue to hold for these problems. It is an interesting problem to compare these methods with other technique for solving quasi variational inclusions for further research activities.

**Keywords:** Convex functions, monotone operators, variational inclusions, iterative methods, inertial methods, convergence.

**AMS 2000 Subject Classification :** 49J40, 90C33

---

\*nkg9607@gmail.com

†Corresponding Author's Email: noormaslam@gmail.com

‡khalidan@gmail.com

§stcsujavarghese@gmail.com

## 1. INTRODUCTION

Quasi variational inclusions, which were introduced and studied by Noor and Noor [28, 30, 35], are useful and important extension of the variational principles with a wide range of applications in industry, physical, regional, social, pure and applied sciences. Quasi variational inclusions provide us with a unified, natural, novel, innovative and general technique to study a wide class of problems arising in different branches of mathematical and engineering sciences, see, for example, [26, 28, 30, 35, 38, 41, 47, 48] and the references therein. One of the most difficult and important problems in variational inequalities is the development of efficient numerical methods. Several numerical methods have been developed for solving the variational inclusions and their variant forms. These methods have been extended and modified in numerous ways. Noor [30, 35] proved that the quasi variational inclusions are equivalent to the fixed point problem. This alternative formulation has allowed to consider the existence of a solution, iterative schemes, sensitivity analysis, merit functions and other aspects of the quasi variational inclusions.

It is very important to develop some efficient iterative methods for solving the quasi variational inequalities. Alvarez et al [1] and Alvarez et al.[2, 3, 4] used the inertial type projection methods for solving variational inequalities. The origin of which can be traced back to Polyak [42]. Noor [32] suggested and investigated inertial type projection methods for solving general variational inequalities. These inertial type methods have been modified in various directions for solving variational inequalities and related optimization problems. Jabeen et al [11, 12, 13] analyzed some inertial projection methods for some classes of general quasi variational inequalities. Convergence analysis of these inertial type methods has been considered under some mild conditions. For more details and applications of the inertial methods, see [1, 2, 3, 4, 9, 10, 11, 12, 13, 32, 36, 40, 41, 42, 43, 47, 48] and reference therein.

Motivated and inspired by the recent research activities, we consider and study some new classes of variational inclusions involving three arbitrary operators, which are called the general quasi variational inclusions. Some special important cases are also discussed. We have shown that the general quasi variational inclusions are equivalent to the fixed point problems. This equivalence is used to study the existence of a solution as well as to propose some new inertial type methods for solving general quasi variational inclusion. The convergence of the proposed inertial methods requires only the Lipschitz continuous of the operators. This fact improves the known convergence criteria of the known results. Note that our method is simple and does not need the strongly

monotonicity of the involved operators. We have only considered theoretical aspects of the suggested methods. It is an interesting problem to implement these methods and to illustrate the efficiency. Comparison with other methods need further research efforts. The ideas and techniques of this paper may be extended for other classes of quasi variational inclusions and related optimization problems.

**2. BASIC DEFINITIONS AND RESULTS**

Let  $H$  be a real Hilbert space whose inner product and norm are denoted by  $\langle \cdot, \cdot \rangle$  and  $\| \cdot \|$  respectively. Let  $\mathcal{A}(\cdot, \cdot) : H \times H \rightarrow H$  be a maximal monotone operator with respect to the first argument. For given nonlinear single valued operator  $\mathcal{T} : H \rightarrow H$  and an arbitrary nonlinear operator  $g : H \rightarrow H$  consider the problem of finding  $\mu \in H$  such that

$$0 \in \rho\mathcal{T}(\mu) + \mu - g(\mu) + \rho\mathcal{A}(\mu, \mu), \tag{2.1}$$

which is called the general quasi variational inclusion, where  $\rho > 0$  is a constant. A number of problems arising in structural analysis, mechanics and economics can be studied in the framework of the general quasi variational inclusions in a unified manner. We now discuss some interesting and important problems, which can be obtained from the problem (2.1).

**Special Cases.**

I. If  $g = I$ , the identity operator, then problem (2.1) reduces to finding  $\mu \in H$  such that

$$0 \in \mathcal{T}(\mu) + \mathcal{A}(\mu, \mu), \tag{2.2}$$

which is called the quasi variational inclusion, introduced and studied by Noor and Noor [35].

II. If  $\mathcal{A}(\cdot, \mu) = \partial\varphi(\cdot, \mu) : H \times H \rightarrow R \cup \{+\infty\}$ , the subdifferential of a convex, proper and lower semi-continuous function  $\varphi(\cdot, \mu)$  with respect to the first argument, then problem (2.1) is equivalent to finding  $\mu \in H$  such that

$$\langle \mathcal{T}(\mu) + \mu - g(\mu), \nu - \mu \rangle + \varphi(\nu, \mu) - \varphi(\mu, \mu) \geq 0, \quad \forall \nu \in H, \tag{2.3}$$

which is called the mixed general quasi variational inequality.

III. If  $\mathcal{A}(g(\mu, \nu)) \equiv \mathcal{A}(\mu)$ , for all  $\nu \in H$ , then problem (2.1) is equivalent to finding  $\mu \in H$  such that

$$0 \in \rho\mathcal{T}(\mu) + \mu - g(\mu) + \rho\mathcal{A}(\mu), \tag{2.4}$$

a problem considered and studied by Noor [28, 30] using the resolvent equations technique.

IV. If  $\mathcal{A}(\mu) \equiv \partial\varphi(\mu)$  is the subdifferential of a proper, convex and lower, semicontinuous function  $\varphi : H \rightarrow R \cup \{+\infty\}$ , then problem (2.4) reduces to: find  $\mu \in H$  such that

$$\langle \mathcal{T}(\mu) + \mu - g(\mu), \nu - \mu \rangle + \varphi(\nu) - \varphi(\mu) \geq 0. \quad (2.5)$$

Problem (2.5) is known as the general mixed variational inequality.

V. If the function  $\varphi(\cdot, \cdot)$  is the indicator function of a closed convex-valued set  $\mathcal{K}(\mu)$  in  $H$ , that is,

$$\varphi(\mu, \mu) = \mathcal{K}_{(\mu)}(\mu) = \begin{cases} 0, & \text{if } \mu \in \mathcal{K}(\mu) \\ +\infty, & \text{otherwise,} \end{cases}$$

then problem (2.3) is equivalent to finding  $\mu \in H$ , such that

$$\langle \mathcal{T}(\mu) + \mu - g(\mu), \nu - \mu \rangle \geq 0, \quad \forall \nu \in \mathcal{K}(\mu), \quad (2.6)$$

a problem considered and studied by Noor et al [40] using the projection method and the implicit Wiener-Hopf equations technique. They have also considered and analyzed some inertial type iterative methods for solving general quasi variational inequalities (2.6). For formulation, motivation, numerical methods, sensitivity analysis, dynamical systems and other aspects of quasi variational inequalities, see [5, 9, 11, 12, 13, 14, 16, 19, 22, 23, 25, 26, 31, 33, 35, ?, ?, 37, 41, 43] and the references therein

VI. If  $\mathcal{K}(\mu) = \mathcal{K}$ , a convex set and  $g = I$ , the identity operator, then then problem (2.6) reduces to finding  $\mu \in \mathcal{K}$  such that

$$\langle \mathcal{T}(\mu), \nu - \mu \rangle \geq 0, \quad \forall \nu \in \mathcal{K}, \quad (2.7)$$

which is called the variational inequality, introduced and studied by Stampacchia [44] in 1964. Variational inequalities can be viewed as novel extensions of the variational principles. Variational inequality has influenced several branches of mathematical, engineering, economics, transforation, regional and medical sciences and continue to inspire researchers to find its applications. For more details, see [5, 6, 7, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48].

VII. If

$$\mathcal{K}^*(\mu) = \{\mu \in H, \langle \mu, \nu \rangle \geq 0, \quad \forall \nu \in \mathcal{K}(\mu)\}$$

is a polar cone of the convex-valued cone  $\mathcal{K}(\mu)$  in  $H$  and  $g = I$ , , then problem (2.6) is equivalent to finding  $\mu \in H$ , such that

$$\mu \in \mathcal{K}(\mu), \quad \mathcal{T}(\mu) \in \mathcal{K}^*(\mu), \quad \langle \mathcal{T}(\mu), \mu \rangle = 0, \quad (2.8)$$

which is called the generalized quasi complementarity problem.

VIII. If  $\mathcal{K}^*(\mu) = \mathcal{K}^*$  then problem (2.8) reduces to finding  $\mu \in \mathcal{H}$  such that

$$\mu \in \mathcal{K}, \quad \mathcal{T}(\mu) \in \mathcal{K}^*, \quad \langle \mathcal{T}(\mu), \mu \rangle = 0, \tag{2.9}$$

which is known as the nonlinear complementarity problem, introduced by Karamardian [15]. Karamardian [15] proved that the nonlinear complementarity problems are equivalent to the variational inequalities. This result enabled us to use the techniques of complementarity problems to solve the variational inequalities and vice versa. For the applications, formulations and generalizations of the complementarity problems, see [7, 15, 32, 37].

**Remark 2.1.** *For special choices of the operators  $\mathcal{T}$ ,  $g$ ,  $\mathcal{A}(\cdot, \cdot)$  and the convex set  $\mathcal{K}$ , one can obtain a large number of implicit (quasi) complementarity problems and variational inequality problems, which are very special cases of problem (2.1). Thus it is clear that problem (2.1) is general and unifying one and has numerous applications in pure and applied sciences.*

We now recall some well known results and notions.

**Definition 2.1 [2].** If  $\mathcal{T}$  is a maximal monotone operator on  $H$ , then, for a constant  $\rho > 0$ , the resolvent operator associated with  $\mathcal{T}$  is defined by

$$\mathcal{J}_{\mathcal{T}}(\mu) = (I + \rho\mathcal{T})^{-1}(\mu), \quad \forall \mu \in H,$$

where  $I$  is the identity operator. Also the resolvent operator  $\mathcal{J}_{\mathcal{T}}$  is single-valued and nonexpansive.

**Remark 2.2.** *Since the operator  $\mathcal{A}(\cdot, \cdot)$  is a maximal monotone operator with respect to the first argument, we denote by*

$$\mathcal{J}_{\mathcal{A}(\mu)} \equiv (I + \rho\mathcal{A}(\mu))^{-1}(\mu), \quad \forall \mu \in H,$$

*the resolvent operator associated with  $\mathcal{A}(\cdot, \mu) \equiv \mathcal{A}(\mu)$ . For example, if  $\mathcal{A}(\cdot, \mu) = \partial\varphi(\cdot, \mu)$ , for all  $\mu \in H$ , and  $\varphi(\cdot, \cdot) : H \times H \rightarrow R \cup \{\infty\}$  is a proper, convex and lower semicontinuous with respect to the first argument, then it is well known that  $\partial\varphi(\cdot, \mu)$  is a maximal monotone operator with respect to the first argument. In this case, the resolvent operator  $\mathcal{J}_{\mathcal{A}(\mu)} = \mathcal{J}_{\varphi(\mu)}$  is defined as*

$$\mathcal{J}_{\varphi(\mu)} = (I + \rho\partial\varphi(\cdot, \mu))^{-1}(\mu) = (I + \rho\partial\varphi(\mu))^{-1}, \quad \forall \mu \in H,$$

*which is defined everywhere on the whole space  $H$ , where  $\partial\varphi(\mu) \equiv \partial\varphi(\cdot, \mu)$ .*

We need the following well-known definitions and results in obtaining our results.

**Definition 2.3.** Let  $T : H \longrightarrow H$  be a given mapping.

*i.* The mapping  $T$  is called  $r$ -strongly monotone ( $r \geq 0$ ), if

$$\langle T\mu - T\nu, \mu - \nu \rangle \geq r \|\mu - \nu\|^2, \quad \forall \mu, \nu \in H.$$

*ii.* The mapping  $T$  is called  $\xi$ -cocoercive ( $\xi > 0$ ), if

$$\langle T\mu - T\nu, \mu - \nu \rangle \geq \xi \|T\mu - T\nu\|^2, \quad \forall \mu, \nu \in H.$$

*iii.* The mapping  $T$  is called relaxed ( $\xi, r$ )-cocoercive ( $r > 0, \xi > 0$ ), if

$$\langle T\mu - T\nu, \mu - \nu \rangle \geq -\xi \|T\mu - T\nu\|^2 + r \|\mu - \nu\|^2, \quad \forall \mu, \nu \in H.$$

For  $\xi = 0$ ,  $T$  is  $r$ -strongly monotone. The class of relaxed ( $\xi, r$ )-cocoercive mapping is the generalized class than the  $r$ -strongly monotone mapping and  $\xi$ -cocoercive.

*iv.* The mapping  $T$  is called  $\eta$ -Lipschitz continuous ( $\eta_1 > 0$ ), if

$$\|T\mu - T\nu\| \leq \eta_1 \|\mu - \nu\|, \quad \forall \mu, \nu \in H.$$

The implicit resolvent operator  $\mathcal{J}_A(\mu)$  is nonexpansive and has the following characterization.

**Assumption [35].** The implicit resolvent operator  $\mathcal{J}_{A(\mu)}$ , satisfies the condition

$$\|\mathcal{J}_{A(\mu)}[\omega] - \mathcal{J}_{A(\nu)}[\omega]\| \leq \eta \|\mu - \nu\|, \quad \forall \mu, \nu, \omega \in H, \quad (2.10)$$

where  $\eta > 0$  is a constant.

**Lemma 2.4.** [42] Consider a sequence of non negative real numbers  $\{\varrho_n\}$ , satisfying

$$\varrho_{n+1} \leq (1 - \Upsilon_n)\varrho_n + \Upsilon_n \sigma_n + \varsigma_n, \quad \forall n \geq 1,$$

where

*i.*  $\{\Upsilon_n\} \subset [0, 1]$ ,  $\sum_{n=1}^{\infty} \Upsilon_n = \infty$ ;

*ii.*  $\limsup \sigma_n \leq 0$ ;

*iii.*  $\varsigma_n \geq 0$  ( $n \geq 1$ ),  $\sum_{n=1}^{\infty} \varsigma_n < \infty$ .

Then,  $\varrho_n \longrightarrow 0$  as  $n \longrightarrow \infty$ .

### 3. MAIN RESULTS

In this section, we prove the equivalence between the problem (2.1) and the fixed point problems. This alternative formulation is used to discuss the existence of a solution as well as to suggest some new inertial-type approximation schemes for solving the general quasi variational inclusion (2.1).

**Lemma 3.1.** *The function  $\mu \in \mathcal{H}$  is a solution of the general quasi variational inclusion (2.1), if and only if,  $\mu \in \mathcal{H}$  satisfies the relation*

$$\mu = \mathcal{J}_{\mathcal{A}(\mu)}[g(\mu) - \rho\mathcal{T}\mu], \tag{3.1}$$

where  $\mathcal{J}_{\mathcal{A}(\mu)}$  is the resolvent operator and  $\rho > 0$  is a constant.

*Proof.* Let  $\mu \in \mathcal{H}$  be a solution of (2.1), then, for a constant  $\rho$ ,

$$\begin{aligned} \rho\mathcal{T}\mu + \mu - g(\mu) + \rho\mathcal{A}(\mu, \mu) &\ni 0, \\ \iff \\ -g(\mu) + \rho\mathcal{T}\mu + \mu + \rho\mathcal{A}(\mu, \mu) &\ni 0 \\ \iff \\ \mu &= \mathcal{J}_{\mathcal{A}}[g(\mu) - \rho\mathcal{T}\mu]. \end{aligned}$$

the required (3.1). □

Lemma 3.1 implies that the general quasi variational inclusion (2.1) is equivalent to the fixed point problem (3.1).

We now define the mapping  $\Phi$  associated with (3.1) as:

$$\Phi(\mu) = \mathcal{J}_{\mathcal{A}(\mu)}[g(\mu) - \rho\mathcal{T}\mu], \tag{3.2}$$

To prove the existence of the solution of problem (2.1), it is enough that the mapping  $\Phi$  defined by (3.2) is a contraction mapping.

**Theorem 3.2.** *Let the operators  $\mathcal{T}, g$  be Lipschitz continuous with constants  $\beta > 0, \sigma > 0$  respectively. If there exists a constant  $\rho > 0$ , such that*

$$\rho < \frac{1 - k}{\beta}, \quad k < 1, \tag{3.3}$$

where

$$k = \sigma + \eta, \tag{3.4}$$

then there exists a solution  $\mu \in \mathcal{H}$  satisfying problem (2.1).

*Proof.* Let  $u \neq v \in H$  be two solutions of problem (2.1). Then, from problem (3.2), we have

$$\begin{aligned}
 \|\Phi(v) - \Phi(u)\| &\leq \|\mathcal{J}_{\mathcal{A}(v)}[g(v) - \rho\mathcal{T}v] - \mathcal{J}_{\mathcal{A}(\mu)}[g(\mu) - \rho\mathcal{T}\mu]\| \\
 &= \|\mathcal{J}_{\mathcal{A}(v)}[g(v) - \rho\mathcal{T}v] - \mathcal{J}_{\mathcal{A}(\mu)}[g(v) - \rho\mathcal{T}v]\| \\
 &\quad + \|\mathcal{J}_{\mathcal{A}(\mu)}[g(v) - \rho\mathcal{T}v] - \mathcal{J}_{\mathcal{A}(\mu)}[g(\mu) - \rho\mathcal{T}\mu]\| \\
 &\leq \|g(v) - g(\mu) - \rho(\mathcal{T}v - \mathcal{T}\mu)\| + \eta\|v - \mu\| \\
 &\leq \|g(v) - g(\mu)\| + \rho\|\mathcal{T}v - \mathcal{T}\mu\| + \eta\|v - \mu\| \\
 &\leq (\sigma + \rho\beta)\|v - \mu\| = \theta\|v - \mu\|, \tag{3.5}
 \end{aligned}$$

since the operators  $\mathcal{T}, g$  are Lipschitz continuous with constant  $\beta > 0$ , and  $\sigma > 0$ , where

$$\theta = \rho\beta + \sigma + \eta = \rho\beta + k \tag{3.6}$$

and  $k$  is defined by (3.4). From (3.3), it follows that  $\theta < 1$ . Thus it follows that the mapping  $\Phi(\mu)$  defined by (3.2) is a contraction mapping and consequently, the mapping  $\Phi(\mu)$  has a fixed point  $\Phi(\mu) = \mu \in \mathcal{H}$  satisfying (2.1), the required result.  $\square$

Using the result (3.1), we can propose some iterative approximation schemes for solving the general quasi variational inclusion (2.1).

**Algorithm 3.1.** For given  $\mu_0 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\mu_{n+1} = (1 - \alpha_n)\mu_n + \alpha_n\{\mathcal{J}_{\mathcal{A}(\mu_n)}[g(\mu_n) - \rho\mathcal{T}\mu_n]\}, \quad n = 1, 2, \dots,$$

where  $\alpha_n \in [0, 1]$ . Algorithm 3.1 is called the Mann iterative resolvent method.

**Algorithm 3.2.** For given  $\mu_0 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\mu_{n+1} = (1 - \alpha_n)\mu_n + \alpha_n\{\mathcal{J}_{\mathcal{A}(\mu_n)}[g(\mu_n) - \rho\mathcal{T}\mu_{n+1}]\}, \quad \alpha_n \in [0, 1]$$

which is known as the extraresolvent method in the sense of Koperlevich [17] and is equivalent to the following two-step method.

**Algorithm 3.3.** For given  $\mu_0 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\begin{aligned}
 \omega_n &= \mathcal{J}_{\mathcal{A}(\mu_n)}[\rho\mathcal{T}\mu_n] \\
 \mu_{n+1} &= (1 - \alpha_n)\mu_n + \alpha_n\{\mathcal{J}_{\mathcal{A}(\mu_n)}[g(\mu_n) - \rho\mathcal{T}\omega_n]\}, \quad \alpha_n \in [0, 1],
 \end{aligned}$$

which is called the predictor-corrector resolvent method.

Using the equation(3.1), we suggest the following double resolvent method:

**Algorithm 3.4.** For given  $\mu_0 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\mu_{n+1} = \mathcal{J}_{A(\mu_n)} [g(\mu_{n+1}) - \rho T\mu_{n+1}], \quad \alpha_n \in [0, 1],$$

Using the predictor-corrector technique, Algorithm 3.4 be written in the following form

**Algorithm 3.5.** For given  $\mu_0 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\begin{aligned} \omega_n &= \mathcal{J}_{A(\mu_n)} [g(\mu_n) - \rho T\mu_n] \\ \mu_{n+1} &= (1 - \alpha_n)\mu_n + \alpha_n \{ \mathcal{J}_{A(\mu_n)} [g(\omega_n) - \rho T\omega_n] \}, \quad \alpha_n \in [0, 1]. \end{aligned}$$

Algorithm 3.5 appeared to be a new two-step method for solving general quasi variational inclusion (2.1).

We can rewrite (3.1 as

$$\mu = \mathcal{J}_{A(\mu)} \{ g((1 - \theta_n)\mu + \theta_n\mu) - \rho T\mu \},$$

where  $\theta_n \in [0, 1], \forall n \geq 1$ .

This fixed point formulation is used to suggest the following two-step method for solving general quasi variational inclusion (2.1) using the ideas of Alvarez [5], Alvarez et al. [6] and Noor [4].

**Algorithm 3.6.** For given  $\mu_0, \mu_1 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\begin{aligned} \omega_n &= \mu_n - \theta_n (\mu_n - \mu_{n-1}) \\ \mu_{n+1} &= 1 - \alpha_n \mu_n + \alpha_n \{ \mathcal{J}_{A(\mu_n)} [\rho T\mu_n] \}, \end{aligned}$$

where  $\theta_n \in [0, 1], \forall n \geq 1$ .

Such type of inertial projection methods for solving general quasi variational inequalities have been considered by Noor [32, 36] and Noor et al[29].

We can rewrite (3.1 as

$$g(\mu) = \mathcal{J}_{A((1-\theta_n)\mu+\theta_n\mu)} \{ g((1 - \theta_n)\mu + \theta_n\mu) - \rho T((1 - \theta_n)\mu + \theta_n\mu) \},$$

where  $\theta_n \in [0, 1], \forall n \geq 1$ .

Using this fixed point formulation, we can suggest the following inertial type methods for solving general quasi variational inclusion (2.1).

**Algorithm 3.7.** For given  $\mu_0, \mu_1 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\begin{aligned} \omega_n &= \mu_n - \theta_n (\mu_n - \mu_{n-1}) \\ g(\mu_{n+1}) &= 1 - \alpha_n \mu_n + \alpha_n \{ \mathcal{J}_{A(\omega_n)} [g(\omega_n) - \rho T\omega_n] \}, \quad n = 1, 2, \dots, \end{aligned}$$

where  $\theta_n \in [0, 1], \forall n \geq 1$ .

Algorithm 3.7 is known as modified inertial resolvent method for solving general quasi variational inclusion (2.1).

**Algorithm 3.8.** For given  $\mu_0, \mu_1 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\begin{aligned}\omega_n &= \mu_n - \theta_n (\mu_n - \mu_{n-1}) \\ \mathbf{g}(\mathbf{y}_n) &= \mathcal{J}_{A(\omega_n)} [\mathbf{g}(\omega_n) - \rho T\omega_n], \\ \mathbf{g}(\mu_{n+1}) &= \mathcal{J}_{A(\mathbf{y}_n)} [\mathbf{g}(\mathbf{y}_n) - \rho T\mathbf{y}_n], \quad n = 1, 2, \dots,\end{aligned}$$

where  $\Theta_n \in [0, 1], \forall n \geq 1$ .

Algorithm 3.8 is a three-step modified inertial method for solving general quasi variational inclusion(2.1).

We now suggest a four-step inertial method for solving the general quasi variational inclusion (2.1).

**Algorithm 3.9.** For given  $\mu_0, \mu_1 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\omega_n = \mu_n - \theta_n (\mu_n - \mu_{n-1}), \quad (3.7)$$

$$\mathbf{x}_n = (1 - \gamma_n)\mu_n + \gamma_n \left\{ \mathcal{J}_{A(\omega_n)} [\mathbf{g}(\omega_n) - \rho T\omega_n] \right\}, \quad (3.8)$$

$$\mathbf{y}_n = (1 - \beta_n)\mu_n + \beta_n \left\{ \mathcal{J}_{A(\mathbf{x}_n)} [\mathbf{g}(\mathbf{x}_n) - \rho T\mathbf{x}_n] \right\}, \quad (3.9)$$

$$\mu_{n+1} = (1 - \alpha_n)\mu_n + \alpha_n \left\{ \mathcal{J}_{A(\mathbf{y}_n)} [\mathbf{g}(\mathbf{y}_n) - \rho T\mathbf{y}_n] \right\}, \quad n = 1, 2, \dots, \quad (3.10)$$

where  $\alpha_n, \beta_n, \gamma_n, \theta_n \in [0, 1], \quad \forall n \geq 1$ .

If  $g = I$ , the identity, then Algorithm (3.9) reduces to:

**Algorithm 3.10.** For given  $\mu_0, \mu_1 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\omega_n = \mu_n - \theta_n (\mu_n - \mu_{n-1}),$$

$$\mathbf{x}_n = (1 - \gamma_n)\mu_n + \gamma_n \left\{ \mathcal{J}_{A(\omega_n)} [(\omega_n) - \rho T\omega_n] \right\},$$

$$\mathbf{y}_n = (1 - \beta_n)\mu_n + \beta_n \left\{ \mathcal{J}_{A(\mathbf{x}_n)} [(\mathbf{x}_n) - \rho T\mathbf{x}_n] \right\},$$

$$\mu_{n+1} = (1 - \alpha_n)\mu_n + \alpha_n \left\{ \mathcal{J}_{A(\mathbf{y}_n)} [(\mathbf{y}_n) - \rho T\mathbf{y}_n] \right\}, \quad n = 1, 2, \dots,$$

where  $\alpha_n, \beta_n, \gamma_n, \theta_n \in [0, 1], \quad \forall n \geq 1$ .

For a different and suitable choice of operators and spaces in Algorithm (3.9), one can obtain numerous new and previous iterative schemes for solving general quasi variational inclusion (2.1) and related problems. This shows that the Algorithm (3.9) is quite flexible and unifying ones. We now estimate convergence analysis for Algorithm 3.9 under some mild and appropriate conditions.

**Theorem 3.3.** *Let the following assumptions be fulfilled:*

- i. The operators  $T, g$  : are  $\eta_1, \eta_2$ -Lipschitz continuous, respectively.*
- ii. Assumption 2 holds.*
- iii. The parameter  $\rho > 0$  satisfies the condition*

$$\rho < \frac{1 - k}{\beta}, \quad k < 1, \tag{3.11}$$

where

$$k = \eta_2 + \nu. \tag{3.12}$$

- iv. Let  $\alpha_n, \beta_n, \gamma_n, \theta_n \in [0, 1]$ , for all  $n \geq 1$  such that*

$$\sum_{n=1}^{\infty} \alpha_n = \infty, \quad \sum_{n=1}^{\infty} \theta_n \|\mu_n - \mu_{n-1}\| < \infty.$$

*Then, for every initial approximation  $\mu_n$ , the sequence  $\{\mu_n\}$  obtained from the iterative scheme defined in Algorithm 3.9 converges to unique solution  $\mu \in H$  satisfying the general quasi variational inclusion (2.1) as  $n \rightarrow \infty$ .*

*Proof.* Let  $\mu \in H : g(\mu) \in K(\mu)$  be a solution of (2.1). Then

$$\mu = (1 - \alpha_n)\mu + \alpha_n \{ \mathcal{J}_{A(\mu)} [g(\mu) - \rho T\mu] \}, \tag{3.13}$$

$$= (1 - \beta_n)\mu + \beta_n \{ \mathcal{J}_{A(\mu)} [g(\mu) - \rho T\mu] \}, \tag{3.14}$$

$$= (1 - \gamma_n)\mu + \gamma_n \{ \mathcal{J}_{A(\mu)} [g(\mu) - \rho T\mu] \}, \tag{3.15}$$

where  $0 \leq \alpha_n, \beta_n, \gamma_n \leq 1, \forall n \geq 1$ , are constants.

Using Assumption (2), from (3.11), and (3.13), we have

$$\begin{aligned} \|\mu_{n+1} - \mu\| &= \|(1 - \alpha_n)\mu_n + \alpha_n \{ \mathcal{J}_{A(y_n)} [g(y_n) - \rho T y_n] \} - (1 - \alpha_n)\mu - \alpha_n \{ \mathcal{J}_{A(\mu)} [g(\mu) - \rho T\mu] \}| \\ &\leq (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n \|\mathcal{J}_{A(y_n)} [g(y_n) - \rho T y_n] - \mathcal{J}_{A(\mu)} [g(\mu) - \rho T\mu]\| \\ &\leq (1 - \alpha_n)\|\mu_n - \mu\| \\ &\quad + \alpha_n \|\mathcal{J}_{A(y_n)} [g(y_n) - \rho T y_n] - \mathcal{J}_{A(y_n)} [g(\mu) - \rho T\mu]\| \\ &\quad + \alpha_n \|\mathcal{J}_{A(y_n)} [g(\mu) - \rho T\mu] - \mathcal{J}_{A(\mu)} [g(\mu) - \rho T\mu]\| \\ &\leq (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n \| [g(y_n) - g(\mu)] - \rho [T y_n - T\mu] \| + \alpha_n \nu \|\mu_n - \mu\| \\ &\leq (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n \|g(y_n) - g(\mu)\| + \alpha_n \rho \| [T y_n - T\mu] \| + \alpha_n \nu \|\mu_n - \mu\|. \\ &\leq (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n (\eta_2 + \alpha_n \rho \eta_1) \|\mu_n - \mu\|, \end{aligned} \tag{3.16}$$

where we have used that fact the operators  $T, g$  are Lischitz continuous with constants  $\eta_1, \eta_2$  From (3.16), we have

$$\begin{aligned}\|\mu_{n+1} - \mu\| &= (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n(\mathbf{k} + \rho\eta_1)\|y_n - \mu\| \\ &= (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n\vartheta\|y_n - \mu\|,\end{aligned}\quad (3.17)$$

where

$$\vartheta = (\mathbf{k} + \eta_1\rho) < 1, \quad \text{from condition (3.11) and } \mathbf{k} \text{ is defined by (3.12).}$$

Similarly, from (3.9) and (3.14), we have

$$\|y_n - \mu\| \leq (1 - \beta_n)\|\mu_n - \mu\| + \beta_n\vartheta\|x_n - \mu\|. \quad (3.18)$$

In a similar way, from (3.11) and (3.18), we have

$$\|x_n - \mu\| \leq (1 - \gamma_n)\|\mu_n - \mu\| + \gamma_n\vartheta\|\omega_n - \mu\|. \quad (3.19)$$

From (3.11), we have

$$\begin{aligned}\|\omega_n - \mu\| &= \|\mu_n - \mu - \theta_n(\mu_n - \mu_{n-1})\| \\ &\leq \|\mu_n - \mu\| + \Theta_n\|\mu_n - \mu_{n-1}\|.\end{aligned}\quad (3.20)$$

From (3.19) and (3.20), we have

$$\begin{aligned}\|x_n - \mu\| &\leq (1 - \gamma_n)\|\mu_n - \mu\| + \gamma_n\vartheta\|\mu_n - \mu\| + \theta_n\|\mu_n - \mu_{n-1}\| \\ &= [1 - \gamma_n(1 - \vartheta)]\|\mu_n - \mu\| + \theta_n\|\mu_n - \mu_{n-1}\| \\ &\leq \|\mu_n - \mu\| + \Theta_n\|\mu_n - \mu_{n-1}\|.\end{aligned}\quad (3.21)$$

From (3.18) and (3.21), we have

$$\begin{aligned}\|y_n - \mu\| &\leq (1 - \beta_n)\|\mu_n - \mu\| + \beta_n\vartheta[\|\mu_n - \mu\| + \theta_n\|\mu_n - \mu_{n-1}\|] \\ &\leq (1 - \beta_n)\|\mu_n - \mu\| + \beta_n\vartheta\|\mu_n - \mu\| + \theta_n\|\mu_n - \mu_{n-1}\| \\ &= [1 - \beta_n(1 - \vartheta)]\|\mu_n - \mu\| + \Theta_n\|\mu_n - \mu_{n-1}\| \\ &\leq \|\mu_n - \mu\| + \Theta_n\|\mu_n - \mu_{n-1}\|.\end{aligned}\quad (3.22)$$

From (3.17) and (3.22), we have

$$\begin{aligned}\|\mu_{n+1} - \mu\| &\leq (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n\vartheta[\|\mu_n - \mu\| + \theta_n\|\mu_n - \mu_{n-1}\|] \\ &\leq (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n\vartheta\|\mu_n - \mu\| + \Theta_n\|\mu_n - \mu_{n-1}\| \\ &= [1 - \alpha_n(1 - \vartheta)]\|\mu_n - \mu\| + \theta_n\|\mu_n - \mu_{n-1}\|.\end{aligned}$$

From condition (3.11), we have  $\vartheta < 1$ . Since  $\sum_{n=1}^{\infty} \alpha_n = \infty$ , setting  $\sigma_n = 0$  and

$$\varsigma_n = \sum_{n=1}^{\infty} \Theta_n \|\mu_n - \mu_{n-1}\| < \infty, \text{ using Lemma 2.4, we have } \mu_n \rightarrow \mu, n \rightarrow \infty.$$

Hence the sequence  $\{\mu_n\}$  obtained from Algorithm 3.9 converges to a unique solution  $\mu \in H$  satisfying the inequality (2.1), the required result.  $\square$

Similarly convergence analysis for other inertial iterative methods can be estimated.

We again suggest some iterative methods by rearranging the equation (3.1). To be more precise, taking  $z = g(\mu) - \rho\mathcal{T}(\mu)$  in (3.1), we have

$$\mu = \mathcal{J}_{\mathcal{A}(\mu)}(z) \tag{3.23}$$

$$z = g(\mu) - \rho\mathcal{T}(\mu) \tag{3.24}$$

$$= \mathcal{J}_{\mathcal{A}(\mu)}(z) - \rho\mathcal{T}(\mu) \tag{3.25}$$

$$= \mathcal{J}_{\mathcal{A}(\mu)}(z) - \rho\mathcal{T}\mathcal{J}_{\mathcal{A}(\mu)}(z). \tag{3.26}$$

Combining (3.24) and (3.25), we obtain

$$g(\mu) = \rho\mathcal{T}(\mu) + \mathcal{J}_{\mathcal{A}(\mu)}(z) - \rho\mathcal{T}\mathcal{J}_{\mathcal{A}(\mu)}(z),$$

which can be written as

$$\mu = (1 - \alpha_n)\mu + \alpha_n\{g(\mu) - \rho\mathcal{T}(\mu) - \mathcal{J}_{\mathcal{A}(\mu)}(z) + \rho\mathcal{T}\mathcal{J}_{\mathcal{A}(\mu)}(z)\}, \tag{3.27}$$

We use the fixed point formulation (3.27) to suggest the following predictor-corrector method for solving the problem (2.1).

**Algorithm 3.11.** For given  $\mu_0, \mu_1 \in H$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\begin{aligned} \omega_n &= \mathcal{J}_{\mathcal{A}(\mu)}[g(\mu_n) - \rho\mathcal{T}(\mu_n)] \\ \mu_{n+1} &= (1 - \alpha_n)\mu_n + \alpha_n\{g(\omega_n) - \rho\mathcal{T}(\omega_n) - g(\nu_n) + \rho\mathcal{T}\nu_n\}, \end{aligned}$$

where  $\alpha_n \in [0, 1]$  is a constant.

The Algorithm 3.11 is known as the forward-backward iterative method.

Now, we again use fixed point formulation (3.27) to suggest the following inertial iterative methods for solving the general quasi variational inclusion (2.1):

**Algorithm 3.12.** For given  $\mu_0, \mu_1 \in \mathbb{H}$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\begin{aligned} g(\omega_n) &= (1 - \beta_n)\mu_n + \beta_n\mu_{n-1} \\ \mu_{n+1} &= (1 - \alpha_n)\mu_n + \alpha_n\{g(\mu_n) - \rho\mathcal{T}(\mu_n) - g(\omega_n) + \rho\mathcal{T}\omega_n\}, \end{aligned}$$

where  $\alpha_n \in [0, 1]$  is a constant.

Algorithm 3.12 is called the forward-backward inertial iterative method for solving the general quasi variational inclusions and this method does not involve any resolvent operator.

From (3.24) and (3.26), we have

$$g(\mu) = \mathcal{J}_{\mathcal{A}(\mu)}[\mathcal{J}_{\mathcal{A}(\mu)}[g(\mu) - \rho\mathcal{T}(\mu)] - \rho\mathcal{T}(\mu)],$$

which can be written as

$$\mu = (1 - \alpha_n)\mu + \alpha_n\{g(\mu) - \mathcal{J}_{\mathcal{A}(\mu)}[\mathcal{J}_{\mathcal{A}(\mu)}[g(\mu) - \rho\mathcal{T}(\mu)] + \rho\mathcal{T}(\mu)]\}, \quad (3.28)$$

where  $\alpha_n \in [0, 1]$  is a constant.

This fixed point formulation is used to propose a new iterative method for solving the general quasi variational inequalities (2.1).

**Algorithm 3.13.** For given  $\mu_0, \mu_1 \in \mathbb{H}$ , compute  $\mu_{n+1}$  by the recurrence relation

$$\mu_{n+1} = (1 - \alpha_n)\mu_n + \alpha_n\{g(\mu_n) - \mathcal{J}_{\mathcal{A}(\mu_n)}[\mathcal{J}_{\mathcal{A}(\mu_n)}[g(\mu_n) - \rho\mathcal{T}(\mu_n)] + \rho\mathcal{T}(\mu_n)]\},$$

where  $\alpha_n \in [0, 1]$  is a constant.

Algorithm 3.13 is called the forward-backward explicit iterative method for solving the general quasi variational inclusions.

#### 4. CONCLUSION:

In this paper, several new inertial projection methods have been suggested and analyzed for solving general quasi variational inclusions. We have proved that the general quasi variational inclusions are equivalent to the implicit fixed point problems. This alternative formulation is used to discuss the existence of a solution of the general quasi variational inclusions. A wide class of three step inertial type iterative methods for solving general quasi variational inclusions are suggested and analyzed. Several

important special cases are discussed. Convergence analysis of these proposed inertial resolvent methods is investigated. It is worth mentioning that Algorithm 3.12 does not involve the calculation of the resolvent operator. We would like to point out that very few numerical examples are available for classical quasi variational inclusions due to their complex nature. In spite of these activities, further efforts are needed to develop numerical implementable methods. It is an interesting problem to compare the efficiency of the proposed methods with other known methods. We expect that the ideas and techniques of this paper will motivate and inspire the interested readers to explore its applications in various fields.

**Data availability:**

Data sharing not applicable to this article as no data sets were generated or analyzed during the current study.

**Conflict interest:**

Authors have no conflict of interest.

**Authors contributions:**

All authors contributed equally to the conception, design of the work, analysis, interpretation of data, reviewing it critically and final approval of the version for publication.

**Acknowledgments:**

The authors wish to express their deepest gratitude to their respected teachers, students, colleagues, collaborators, referees and friends, who have directly or indirectly contributed in the preparation of this paper.

**REFERENCES**

- [1] F. Alvarez, Weak convergence of a relaxed and inertial hybrid projection-proximal point algorithm for maximal monotone operators in Hilbert space, *SIAM J. Optim.* 14(2003), 773-782.
- [2] F. Alvarez and H. Attouch, An inertial proximal method for maximal monotone operators via discretization of a nonlinear oscillator with damping, *Set-Valued Anal.* 9(2001), 3-11.
- [3] H. Attouch and M. O. Czarnecki, Asymptotic control and stabilization of nonlinear oscillators with non-isolated equilibria, *J. Diff. Equat.* 179 (2002), 278-31.

- [4] H. Attouch, X. Goudon and P. Redont, The heavy ball with friction.I. The continuous dynamical system, *Commun. Contemp. Math.* **2**(2000), 1-34 .
- [5] A. Bensoussan and J. L. Lions, *Application des Inequalities Variationnelles en Control eten Stochastique*, Dunod, Paris, (1978).
- [6] A. Bnouhachem and M. A. Noor, Three-step projection method for general variational inequalities, *Int. J. Mod. Phys. B.* **26**(14)(2012).
- [7] R. W. Cottle, Jong-S. Pang and R. E. Stone, *The Linear Complementarity Problem*, SIAM Publication, 1992.
- [8] V. M. Filippov, *Variational Principles for Nonpotential Operators*, Amer. Math. Soc. Providence, Rode Island, USA, 1989.
- [9] A. Dixit, D.R. Sahu, P. Gautam, T. Som and J.C. Yao, An accelerated forward-backward splitting algorithm for solving inclusion problems with applications to regression and link prediction problems, *J. Nonlin. Var. Anal.* **5**(2021), 79-101.
- [10] R. Glowinski, J. L. Lions and R. Tremolieres, *Numerical Analysis of Variational Inequalities*, North-Holland, Amsterdam, 1981.
- [11] S. Jabeen, M. A. Noor and K. I. Noor, Some new inertial projection methods for quasi variational Inequalities, *Appl. Math. E-Notes*, **21**(2021), 172-178.
- [12] S. Jabeen, M. A. Noor and K. I. Noor, Inertial iterative methods for general quasi variational inequalities and dynamical systems, *J. Math. Anal.* **11**(3)(2020), 14-29.
- [13] S. Jabeen, B. Bin- Mohsen, M. A. Noor and K. I. Noor, Inertial projection methods for solving general quasi-variational inequalities, *AIMS Math.* **6**(2)(2021), 1075-1086. doi: 10.3934/math.2021064.
- [14] M. Jacimovic and N. Mijajlovic, On a continuous gradient-type method for solving quasi variational inequalities, *Proc. Montenegrin Acad. Sci Arts.* **19**(2011), 16-27.
- [15] S. Karamardian, Generalized complementarity problem, *J. Optim Theory Appl.* **8**(1971), 161-168.
- [16] D. Kinderlehrer and G. Stampachia, *An Introduction to Variational Inequalities and Their Applications*, . Academic Press, New York, USA, 1980.
- [17] G. M. Koperlevich, The extragradient method for finding saddle points and other problems, *Matecom.*, **12**(1976), 203-213.
- [18] P. E. Mainge, Regularized and inertial algorithms for common fixed points of nonlinear operators, *J. Math. Anal. Appl.* **344**(2008), 876-887
- [19] N. Mijajlovic, J. Milojica and M. A. Noor, Gradient-type projection methods for quasi variational inequalities, *Optim. Letters.* **13**(2019), 1885-1896.

- [20] K. G. Murty: Linear Complementarity, Linear and Nonlinear Programming, Heldermann Verlag, Berlin, 1988.
- [21] M. A. Noor, On Variational Inequalities, Ph.D Thesis, Brunel University, London, UK, 1975.
- [22] M. A. Noor, An iterative scheme for class of quasi variational inequalities, J. Math. Anal. Appl. 110(1985), 463-468.
- [23] M. A. Noor, Noor, General quasi complementarity problems, J. Math. Anal. Appl. 120(1986). 321-327,
- [24] M. A. Noor, General variational inequalities, Appl. Math. Letters, 1(1988), 119-121.
- [25] M. A. Noor, Quasi variational inequalities, Appl. Math. Letters, 1(4)(1988),367-370.
- [26] M. A. Noor, Generalized multivalued quasi-variational inequalities (II), Computer Math. Appl. 35(5)(1988), 63-78
- [27] M. A. Noor, Variational inequalities in physical oceanography, in: Ocean Wave Engineering(Edit. M. Rahman ), Computational Mechanics Publications, Southampton, UK, 1994, 201-226.
- [28] M. A. Noor, Generalized set-valued variational inclusions and resolvent equations, J. Math. Anal. Appl. 228(1998), 206-220.
- [29] M. A. Noor, New approximation schemes for general variational inequalities, J. Math. Anal. Appl. 251(2000), 217-229.
- [30] M. A. Noor, Three-step iterative algorithms for multivalued quasi variational inclusions, J. Math. Anal. Appl. 255(2001), 589-604.
- [31] M. A. Noor, Implicit dynamical systems and quasi variational inequalities, Appl. Math. Comput. 134(2003),69-81.
- [32] M. A. Noor, Some developments in general variational inequalities, Appl. Math. Comput., 152(2004), 199-277.
- [33] M. A. Noor, Merit functions for quasi variational inequalities, J. Math. Inequal. 1(2007), 259-269.
- [34] M. A. Noor, Extended general variational inequalities, Appl. Math. Lett.,22(2009), 182-185.
- [35] M. A. Noor and K. I. Noor, Sensitivity analysis of quasi variational inclusions, J. Math. Anal. Appl. 236(1999), 290-299.

- [36] M. A. Noor, K. I. Noor and M. Th. Rassias, Some new trends in mixed variational inequalities, *J. Advanc. Math. Studies*, 15(2)(2022), 105-140.
- [37] M. A. Noor and W. Oettli, On general nonlinear complementarity problems and quasi equilibria, *Le Mathemath(Catania)*. 49 (1994), 313-331.
- [38] M. A. Noor, K. I. Noor and M. Th. Rassias, New trends in general variational inequalities, *Acta Appl. Mathematica*, 170(1)(2020), 981-1064.
- [39] M. A. Noor, K. I. Noor and T. M. Rassias, Some aspects of variational inequalities, *J. Comput. Appl. Math.* 47(1993), 285-312.
- [40] M. A. Noor, K. I. Noor and B. B. Mohsin, Some new classes of general quasi variational inequalities, *AIMS Math.* 6(6)(2021), 6406-6421.
- [41] M. A. Noor, K. I. Noor, S. Treanta and K. Nonlaopon, On three-step iterative schemes associated with general quasi-variational inclusions, *Alexandria Eng. J.*, 61(2022),12051-12059
- [42] B. T. Polyak, Some methods of speeding up the convergence of iterative methods, *Zh. Vychisl. Mat. Mat. Fiz.* 4(1964), 791-803.
- [43] Y. Shehu, A. Gibali and S. Sagratella, Inertial projection-type method for solving quasi variational inequalities in real Hilbert space, *J. Optim. Theory Appl.* 184(2020), 877-894.
- [44] G. Stampacchia, Formes bilineaires coercivites sur les ensembles convexes, *C. R. Acad. Sci.Paris.*, 258(1964), 4413-4416.
- [45] E. Tonti, Variational formulation for every nonlinear problem, *Intern. J. Eng. Sci.* 22(11-12)(1984), 1343- 1371.
- [46] H. K. Xu, Iterative algorithms for nonlinear operators, *J. Lond. Math. Soc.* 66(2002), 240-256.
- [47] Y. Yao, Y. Shehu, X. Li and Q. Dong, A method with inertial extrapolation step for split monotone inclusion problems, *Optimization*, 70(2021), 741-761.
- [48] X. Zhao, J.C. Yao and Y. Yao, A proximal algorithm for solving split monotone variational inclusions, *U. P. B. Scient, Bulletin-Series A*, 82(3)(2020), 43-52.