

Quantum Algorithm for 3-SAT Problem of 15, and 16 Variables by Quantum Fourier Transform with Repeat Qubits, and Weight Changes on QCEngine

Toru Fujimura

*Art and Physical Education area security office, University of Tsukuba,
Ibaraki-branch, Rising Sun Security Service Co., Ltd., 1-1-1, Tennodai, Tsukuba,
Ibaraki 305-8577, Japan
E-mail: tfujimura8@gmail.com*

Abstract

A quantum algorithm for the 3-SAT problem of 15, and 16 variables by the quantum Fourier transform with the repeat qubits, and the weight changes on the QCEngine, and its example are reported. When there are 3 literals with 2 ‘OR’s in each clause, a weight of r -th clause is m_r [m_r is a natural number. $1 \leq r \leq (\text{number of clauses}) = d$.], the r -th clause is $C_{u,r}(x_1, x_2, x_3, \dots, x_n)$ [u is $2^0x_1 + 2^1x_2 + 2^2x_3 + \dots + 2^{n-1}x_n$. $x_1, x_2, x_3, \dots, x_n$ are the variables, and the repeat qubits.], and $S(u)$ is $\sum_{r=1 \rightarrow d} m_r \times C_{u,r}(x_1, x_2, x_3, \dots, x_n)$, $\text{mod}(S(u)_{\max})$ of $S(u)$ [$S(u)_{\max}$ is the maximum value of $S(u)$.] is computed, next, for u , the quantum Fourier transform is done. In this time, there are 15, and 16 variables, the repeat qubits, and changed weights. The complexity of this method is able to be several times.

Keywords: Quantum algorithm, 3-SAT problem, 15, and 16 variables, quantum Fourier transform, repeat qubits, weight changes, QCEngine.

AMS subject classification: Primary 81-08; Secondary 81-10, 68Q12.

1. Introduction

Deutsch, and Jozsa reported the first quantum algorithm for the exponential accelerated computation. [1] Shor discussed the valid quantum algorithm for factorization. [2] Grover reported the fast quantum algorithm for database search. [3, 4] The complexity of the 3-SAT problem had been discussed by Cook. [5] The quantum computer’s example of the 3-SAT problem is reported by Johnston, Harrigan, and Gimeno-Segovia with the QCEngine (free on-line quantum computer

simulator). [6] Fujimura discussed a quantum algorithm for the 3-SAT problem by the Shor's Fourier transform with the RAM on the QCEngine. [7] Still more, Fujimura discussed a quantum algorithm for the 3-SAT problem of 14 variables by the quantum Fourier transform with the repeat qubits, and the weight changes on the QCEngine. [8]

According to my advanced study, when the quantum Fourier transform with the repeat qubits, and the weight changes on the QCEngine for the 3-SAT problem of 15, and 16 variables is used, the complexity of the 3-SAT problem of 15, and 16 variables is able to be several times.

Therefore, the quantum algorithm for the 3-SAT problem of 15, and 16 variables is examined by the quantum Fourier transform with the repeat qubits, and the weight changes on the QCEngine, and its result are reported.

2. 3-SAT Problem

In the 3-SAT problem, it is assumed that (i) each value of n variables becomes "TRUE", or "FALSE", " \sim " is "NOT", " \vee " is "OR", " $\&$ " is "AND", (ii) " \vee ", " \sim ", and 3 different variables are included in each parentheses (= clause) that are connected by " $\&$ ". If a value of logical formula by the literals, and the logical connectives is "TRUE", it is decided whether there is at least one combination of values of the variables or not. [5-8]

3. Quantum Algorithm

The following conditions are assumed. (I) Each value of variables x_1, x_2, x_3, \dots , and x_n becomes "TRUE" [= 1], or "FALSE" [= 0]. " \sim " is "NOT". " \vee " is "OR". " $\&$ " is "AND". For example, it is assumed in this algorithm that (1 \vee 1 \vee 1), (1 \vee 1 \vee 0), and (1 \vee 0 \vee 0) become 1, and (0 \vee 0 \vee 0) becomes 0. (II) " \vee ", " \sim ", and 3 different variables in x_1, x_2, x_3, \dots , and x_n are included in each clause, and then the clauses are connected by " $\&$ ". In these conditions, if a value of logical formula by the literals, and the operators is "TRUE", it is searched whether there is at least one combination of values of the variables or not. It is assumed that n is number of qubits, u is $2^0x_1 + 2^1x_2 + 2^2x_3 + \dots + 2^{n-1}x_n$, a weight of r -th clause is m_r [m_r is a natural number. $1 \leq r \leq$ (number of clauses) = d .], the r -th clause is $C_{u,r}(x_1, x_2, x_3, \dots, x_n)$, and $S(u)$ is $\sum_{r=1}^d m_r \times C_{u,r}(x_1, x_2, x_3, \dots, x_n)$, and $S(u)_{\max}$ is [the maximum value of $S(u)$] = k .

First of all, query quantum registers $|x_i\rangle$ [$1 \leq i \leq n$. i is an integer. n is the number of variables in the logical formula, and the repeat qubits.], and work1 quantum registers $|w_{1,j}\rangle$ [$1 \leq j \leq t$. j , and t are integers. t is a necessary number for $S(u)_{\max} \leq \sum_{f=0}^t 2^f$. f is an integer.].

Step 1: The r data are introduced to the RAM [6].

Step 2: Each qubit of $|x_i\rangle$, and $|w_{1,j}\rangle$ is set $|0\rangle$.

Step 3: The Hadamard gate \boxed{H} [2-4, 6-10] acts on each qubit of $|x_i\rangle$. It changes them for entangled states.

Step 4: Each clause is presented by $|x_i\rangle$, $|w_{1,j}\rangle$, add gate, and quantum operators. For $|x_i\rangle$, RAM[$r - 1$] [RAM has m_r data of $r = 1 \rightarrow d$.] is incremented in $|w_{1,j}\rangle$. In a function, $S(u) = \sum_{r=1}^d m_r \times C_{u,r}(x_1, x_2, x_3, \dots, x_n)$ is computed. This operation

makes entangled data base. In this case, n is 15, and 16 variables in the logical formula, and the number of repeat qubits.

Step 5: For $|x_i\rangle$, the quantum Fourier transform (= QFT) [2, 6-10] is done.

Step 6: For $|x_i\rangle$, and $|w_{1,j}\rangle$, the probes are done.

Step 7: For $|x_i\rangle$, the read is done.

Step 8: A number of spikes is estimated by the function (<https://oreilly-qc.github.io/?p=12-4> [6]), where the function estimate_num_spikes (spike, range) [spike: read value, range: 2^n] is used.

Step 9: From candidates of the number of spikes, the repeat period P is obtained.

Step 10: From $u = P = 2^0x_1 + 2^1x_2 + 2^2x_3 + \dots + 2^{n-1}x_n$, when there is $S(P)_{\max} [= \sum_{r=1}^d m_r \times C_{P,r}(x_1, x_2, x_3, \dots, x_n) = k]$, it is the answer [one combination of (value of logical formula) = 1].

Step 11: When the probability is 24%, or more than 24%, the next weight changed computation isn't done.

4. Example of Numerical Computation

4-1. 16 Variables

4-1-1. 16 Variables, 5 Repeat Qubits, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]

For example of $n = 21$ [16 variables, and 5 repeat qubits], it is assumed that a logical formula : $(x_3 \vee x_4 \vee x_5) \& (\sim x_1 \vee x_2 \vee x_3) \& (\sim x_3 \vee x_4 \vee x_5) \& (x_3 \vee \sim x_4 \vee x_5) \& (\sim x_2 \vee x_3 \vee \sim x_5) \& (\sim x_3 \vee \sim x_4 \vee x_5) \& (\sim x_3 \vee x_4 \vee \sim x_5) \& (x_3 \vee \sim x_4 \vee \sim x_5) \& (\sim x_3 \vee \sim x_4 \vee \sim x_5) \& (x_4 \vee \sim x_5 \vee \sim x_6) \& (\sim x_5 \vee x_6 \vee \sim x_7) \& (x_6 \vee x_7 \vee \sim x_8) \& (x_7 \vee x_8 \vee \sim x_9) \& (x_8 \vee x_9 \vee \sim x_{10}) \& (x_9 \vee x_{10} \vee \sim x_{11}) \& (x_{10} \vee x_{11} \vee \sim x_{12}) \& (x_{11} \vee x_{12} \vee \sim x_{13}) \& (x_{12} \vee x_{13} \vee \sim x_{14}) \& (x_{13} \vee x_{14} \vee \sim x_{15}) \& (x_{14} \vee x_{15} \vee \sim x_{16})$, each value of x_{1-16} : $x_1 = x_2 = x_3 = x_4 = x_6 = x_7 = x_8 = x_9 = x_{10} = x_{11} = x_{12} = x_{13} = x_{14} = x_{15} = x_{16} = 0$, $x_5 = 1$, $m_1 = 1$, $m_2 = 2$, $m_3 = 3$, $m_4 = 4$, $m_5 = 5$, $m_6 = 6$, $m_7 = 7$, $m_8 = 8$, $m_9 = 9$, $m_{10} = 10$, $m_{11} = 11$, $m_{12} = 12$, $m_{13} = 13$, $m_{14} = 14$, $m_{15} = 15$, $m_{16} = 16$, $m_{17} = 17$, $m_{18} = 18$, $m_{19} = 19$, $m_{20} = 20$, $t = 7$, and $k = (20 + 1)20/2 = 210$. It is noticed that the maximum value of $(n + t)$ on the QCEngine is 28.

An example of program on the QCEngine is the following.

```
10 var a = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]; // RAM_a
20 var query_qubits = 21;
30 var work1_qubits = 7;
40 qc.reset(query_qubits + work1_qubits);
50 var query = qint.new(query_qubits, 'query');
60 var work1 = qint.new(work1_qubits, 'work1');
70 qc.label('q'); // set query
80 query.write(0);
90 query.hadamard();
100 qc.label(' ');
110 qc.label('w1'); // set work1
120 work1.write(0);
130 qc.print(' RAM before increment : ' + a + '\n');
140 var query16 = 16;
```

```
150 var work1_0 = 0;
160 qc.label('increment');
170 qc.not(query.bits(0x4)|query.bits(0x8)|query.bits(0x10));
180 work1.add(a[0],query.bits(0x4)|query.bits(0x8)|query.bits(0x10));
190 qc.not(query.bits(0x4)|query.bits(0x8)|query.bits(0x10));
200 qc.not(query.bits(0x2)|query.bits(0x4));
210 work1.add(a[1],query.bits(0x1)|query.bits(0x2)|query.bits(0x4));
220 qc.not(query.bits(0x2)|query.bits(0x4));
230 qc.not(query.bits(0x8)|query.bits(0x10));
240 work1.add(a[2],query.bits(0x4)|query.bits(0x8)|query.bits(0x10));
250 qc.not(query.bits(0x8)|query.bits(0x10));
260 qc.not(query.bits(0x4)|query.bits(0x10));
270 work1.add(a[3],query.bits(0x4)|query.bits(0x8)|query.bits(0x10));
280 qc.not(query.bits(0x4)|query.bits(0x10));
290 qc.not(query.bits(0x4));
300 work1.add(a[4],query.bits(0x2)|query.bits(0x4)|query.bits(0x10));
310 qc.not(query.bits(0x4));
320 qc.not(query.bits(0x10));
330 work1.add(a[5],query.bits(0x4)|query.bits(0x8)|query.bits(0x10));
340 qc.not(query.bits(0x10));
350 qc.not(query.bits(0x8));
360 work1.add(a[6],query.bits(0x4)|query.bits(0x8)|query.bits(0x10));
370 qc.not(query.bits(0x8));
380 qc.not(query.bits(0x4));
390 work1.add(a[7],query.bits(0x4)|query.bits(0x8)|query.bits(0x10));
400 qc.not(query.bits(0x4));
410 work1.add(a[8],query.bits(0x4)|query.bits(0x8)|query.bits(0x10));
420 qc.not(query.bits(0x8));
430 work1.add(a[9],query.bits(0x8)|query.bits(0x10)|query.bits(0x20));
440 qc.not(query.bits(0x8));
450 qc.not(query.bits(0x20));
460 work1.add(a[10],query.bits(0x10)|query.bits(0x20)|query.bits(0x40));
470 qc.not(query.bits(0x20));
480 qc.not(query.bits(0x20)|query.bits(0x40));
490 work1.add(a[11],query.bits(0x20)|query.bits(0x40)|query.bits(0x80));
500 qc.not(query.bits(0x20)|query.bits(0x40));
510 qc.not(query.bits(0x40)|query.bits(0x80));
520 work1.add(a[12],query.bits(0x40)|query.bits(0x80)|query.bits(0x100));
530 qc.not(query.bits(0x40)|query.bits(0x80));
540 qc.not(query.bits(0x80)|query.bits(0x100));
550 work1.add(a[13],query.bits(0x80)|query.bits(0x100)|query.bits(0x200));
560 qc.not(query.bits(0x80)|query.bits(0x100));
570 qc.not(query.bits(0x100)|query.bits(0x200));
580 work1.add(a[14],query.bits(0x100)|query.bits(0x200)|query.bits(0x400));
590 qc.not(query.bits(0x100)|query.bits(0x200));
```

```

600 qc.not(query.bits(0x200)|query.bits(0x400));
610 work1.add(a[15],query.bits(0x200)|query.bits(0x400)|query.bits(0x800));
620 qc.not(query.bits(0x200)|query.bits(0x400));
630 qc.not(query.bits(0x400)|query.bits(0x800));
640 work1.add(a[16],query.bits(0x400)|query.bits(0x800)|query.bits(0x1000));
650 qc.not(query.bits(0x400)|query.bits(0x800));
660 qc.not(query.bits(0x800)|query.bits(0x1000));
670 work1.add(a[17],query.bits(0x800)|query.bits(0x1000)|query.bits(0x2000));
680 qc.not(query.bits(0x800)|query.bits(0x1000));
690 qc.not(query.bits(0x1000)|query.bits(0x2000));
700 work1.add(a[18],query.bits(0x1000)|query.bits(0x2000)|query.bits(0x4000));
710 qc.not(query.bits(0x1000)|query.bits(0x2000));
720 qc.not(query.bits(0x2000)|query.bits(0x4000));
730 work1.add(a[19],query.bits(0x2000)|query.bits(0x4000)|query.bits(0x8000));
740 qc.not(query.bits(0x2000)|query.bits(0x4000));
750 qc.label('QFT');
760 query.QFT();
770 var prob16 = 0;
780 prob16 += query.peekProbability(query16);
790 // Print output query-Prob
800 qc.print(' Prob_query16: ' + prob16);
810 var prob0 = 0;
820 prob0 += work1.peekProbability(work1_0);
830 // Print output work1-Prob
840 qc.print(' Prob_work1_0: ' + prob0);
850 //read
860 qc.label('Rq');
870 var b2 = query.read();
880 // Print output result
890 qc.print(' Read query = ' + b2 + '.');
900 // end

```

When this program is copied on Programming Quantum Computers <https://oreilly-qc.github.io/#> [free on-line quantum computation simulator QCEngine] [6], you can run it. [Caution!: Please delate the line numbers.]

A result of this program is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000015259 (\approx 1/65536)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.000000000$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 2076800, 1808384, 1935360, 1118272, 4352, 305024, 1261824, 1712640, 1992192, 1531872, 2028352, 229312, 754176, 46560, 0, 1671168, 0, 1852096, 590848, 2004992, 1992512, 1042688, 2037760, 1352960, 66560, 1500800, 1509376, 197632, 64512, 1539680, 819840, 2095104, 130944, 393472, 884864, 1999872, 122912, 853056, 81920, 752128, 724480, 1368576, 345088, 1787872, 128, 441248, 1636864, 1497344, 921088, 775424. (= spike)$

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{21} = 2099152$]] : $R_q \rightarrow$ candidates ; 2076800 \rightarrow 103, ... ; 1808384 \rightarrow 7, 15, 22, ... ; 1935360 \rightarrow 13, 26, ... ; 1118272 \rightarrow 2, 4, 6, 9, 11, 13, 15, 30, ... ; 4352 \rightarrow 482, ... ; 305024 \rightarrow 7, 14, 21, ... ; 1261824 \rightarrow 3, 5, 10, 15, 20, ... ; 1712640 \rightarrow 5, 11, 22, ... ; 1992192 \rightarrow 20, ... ; 1531872 \rightarrow 4, 7, 11, 15, 26, ... ; 2028352 \rightarrow 30, ... ; 229312 \rightarrow 9, 18, ... ; 754176 \rightarrow 3, 6, 8, 11, 14, 25, ... ; 46560 \rightarrow 45, ... ; 0 \rightarrow nothingness ; 1671168 \rightarrow 5, 10, 15, 20, ... ; 1852096 \rightarrow 9, 17, ... ; 590848 \rightarrow 4, 7, 14, 18, ... ; 2004992 \rightarrow 9, 17, ... ; 1992512 \rightarrow 20, ... ; 1042688 \rightarrow 2, 4, 6, 8, 10, 12, 14, 16, ... ; 2037760 \rightarrow 35, ... ; 1352960 \rightarrow 3, 6, 8, 11, 14, 17, ... ; 66560 \rightarrow 32, ... ; 1500800 \rightarrow 4, 7, 14, 21, ... ; 1509376 \rightarrow 4, 7, 14, 18, ... ; 197632 \rightarrow 11, 21, ... ; 64512 \rightarrow 33, ... ; 1539680 \rightarrow 4, 8, 11, 15, 30, ... ; 819840 \rightarrow 3, 5, 10, 13, 18, ... ; 2095104 \rightarrow 1024, ... ; 130944 \rightarrow 16, ... ; 393472 \rightarrow 5, 11, 16, ... ; 884864 \rightarrow 2, 5, 7, 12, 19, ... ; 1999872 \rightarrow 22, ... ; 122912 \rightarrow 17, ... ; 853056 \rightarrow 3, 5, 10, 15, 17, ... ; 81920 \rightarrow 26, ... ; 752128 \rightarrow 3, 6, 8, 11, 14, 25, ... ; 724480 \rightarrow 3, 6, 9, 12, 14, 17, ... ; 1368576 \rightarrow 3, 6, 9, 12, 14, 17, ... ; 345088 \rightarrow 6, 12, 18, ... ; 1787872 \rightarrow 7, 14, 20, ... ; 128 \rightarrow 16384, ... ; 441248 \rightarrow 5, 10, 14, 19, ... ; 1636864 \rightarrow 5, 9, 18, ... ; 1497344 \rightarrow 4, 7, 14, 21, ... ; 921088 \rightarrow 2, 5, 7, 9, 16, ... ; 775424 \rightarrow 3, 5, 8, 16,

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} + 2^{16}x_{17} + 2^{17}x_{18} + 2^{18}x_{19} + 2^{19}x_{20} + 2^{20}x_{21} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 + 2^{16} \times 0 + 2^{17} \times 0 + 2^{18} \times 0 + 2^{19} \times 0 + 2^{20} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-1-2. 16 Variables, 4 Repeat Qubits, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]

For example of $n = 20$ [16 variables, and 4 repeat qubits], it is assumed that the logical formula, each value of $x_{1\sim 16}$, each value of $m_{1\sim 20}$, t , and k are same in the section 4-1-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000015259 (\approx 1/65536)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.00081668$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 395168, 361984, 8208, 231424, 241984, 34240, 32576, 1016816, 964160, 1014272, 45184, 55296, 363648, 32832, 167744, 848640, 978944, 984224, 1016384, 1020032, 915968, 162432, 20160, 159744, 36864, 416832, 458752, 132160, 884480, 31232, 20480, 950880, 340992, 990720, 97984, 951360, 0, 966528, 818176, 622336, 98272, 88544, 1007600, 754688, 0, 322656, 687936, 990848, 41600, 26624. (= spike)$

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{20} = 1048576$]] : $R_q \rightarrow$ candidates ; 395168 \rightarrow 3, 5, 8, 16, ... ; 361984 \rightarrow 3, 6, 9, 12, 15, 17, ... ; 8208 \rightarrow 128, ... ; 231424 \rightarrow 5, 9, 18, ... ; 241984 \rightarrow 4, 9, 13, 26, ... ; 34240 \rightarrow 31, ... ; 32576 \rightarrow 32, ... ; 1016816 \rightarrow 33, ... ; 964160 \rightarrow 12, 25, ... ; 1014272 \rightarrow 31, ... ; 45184 \rightarrow 23, ... ; 55296 \rightarrow 19, ... ; 363648 \rightarrow 3, 6, 9, 12, 14, 17, ... ; 32832 \rightarrow 32, ... ; 167744 \rightarrow 6, 13, 19, ... ; 848640 \rightarrow 5, 11, 16, ... ; 978944 \rightarrow 15, 30, ... ; 984224 \rightarrow

16, ... ; 1016384 → 33, ... ; 1020032 → 37, ... ; 915968 → 8, 16, ... ; 162432 → 6, 13, 26, ... ; 20160 → 52, ... ; 159744 → 7, 13, 26, ... ; 36864 → 28, ... ; 416832 → 3, 5, 10, 15, 20, ... ; 458752 → 2, 5, 7, 9, 16, ... ; 132160 → 8, 16, ... ; 884480 → 6, 13, 19, ... ; 31232 → 34, ... ; 20480 → 51, ... ; 950880 → 11, 21, ... ; 340992 → 3, 6, 9, 12, 15, 18, ... ; 990720 → 18, ... ; 97984 → 11, 21, ... ; 951360 → 11, 22, ... ; 0 → nothingness ; 966528 → 13, 26, ... ; 818176 → 5, 9, 18, ... ; 622336 → 3, 5, 10, 15, 17, ... ; 98272 → 11, 21, ... ; 88544 → 12, 24, ... ; 1007600 → 26, ... ; 754688 → 4, 7, 14, 18, ... ; 322656 → 3, 7, 10, 13, 26, ... ; 687936 → 3, 6, 9, 12, 15, 17, ... ; 990848 → 18, ... ; 41600 → 25, ... ; 26624 → 39,

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} + 2^{16}x_{17} + 2^{17}x_{18} + 2^{18}x_{19} + 2^{19}x_{20} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 + 2^{16} \times 0 + 2^{17} \times 0 + 2^{18} \times 0 + 2^{19} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-1-3. 16 Variables, 3 Repeat Qubits, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]

For example of $n = 19$ [16 variables, and 3 repeat qubits], it is assumed that the logical formula, each value of $x_{1\sim 16}$, each value of $m_{1\sim 20}$, t , and k are same in the section 4-1-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000015259 (\approx 1/65536)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.00041169$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 14592, 233472, 147184, 516352, 411136, 16320, 153408, 449408, 487440, 507520, 179232, 345184, 14848, 344608, 129984, 105792, 508056, 172032, 81616, 478976, 47152, 8256, 497728, 523904, 469056, 0, 170080, 468480, 8192, 432992, 511504, 65072, 57408, 32640, 516648, 49368, 256096, 146464, 487680, 473608, 32768, 516096, 315488, 182272, 494784, 233472, 33192, 264128, 425472, 48128. (= spike)$

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{19} = 524288$]] : $R_q \rightarrow$ candidates ; 14592 → 36, ... ; 233472 → 2, 5, 7, 9, 18, ... ; 147184 → 4, 7, 14, 18, ... ; 516352 → 66, ... ; 411136 → 5, 9, 14, 28, ... ; 16320 → 32, ... ; 153408 → 3, 7, 14, 17, ... ; 449408 → 7, 14, 21, ... ; 487440 → 14, 28, ... ; 507520 → 31, ... ; 179232 → 3, 6, 9, 12, 15, 18, ... ; 345184 → 3, 6, 9, 12, 15, 18, ... ; 14848 → 35, ... ; 344608 → 3, 6, 9, 12, 15, 18, ... ; 129984 → 4, 8, 12, 16, ... ; 105792 → 5, 10, 15, 20, ... ; 508056 → 32, ... ; 172032 → 3, 6, 9, 12, 15, 18, ... ; 81616 → 6, 13, 26, ... ; 478976 → 12, 23, ... ; 47152 → 11, 22, ... ; 8256 → 64, ... ; 497728 → 20, ... ; 523704 → 1365, ... ; 469056 → 10, 19, ... ; 0 → nothingness ; 170080 → 3, 6, 9, 12, 15, 19, ... ; 468480 → 9, 19, ... ; 8192 → 64, ... ; 432992 → 6, 12, 17, ... ; 511504 → 41, ... ; 65072 → 8, 16, ... ; 57408 → 9, 18, ... ; 32640 → 16, ... ; 516648 → 69, ... ; 49368 → 11, 21, ... ; 256096 → 2, 4, 6, 8, 10, 12, 14, 16, ... ; 146464 → 4, 7, 14, 18, ... ; 487680 → 14, 29, ... ; 473608 → 10, 21, ... ; 32768 → 16, ... ; 516096 → 64, ... ; 315488 → 3, 5, 10, 15, 20, ... ; 182272 → 3, 6, 9, 12, 14, 17, ... ; 494784 →

18, ... ; 233472 → 2, 5, 7, 9, 18, ... ; 33192 → 16, ... ; 264128 → 2, 4, 6, 8, 10, 12, 14, 16, ... ; 425472 → 5, 11, 16, ... ; 48128 → 11, 22,

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} + 2^{16}x_{17} + 2^{17}x_{18} + 2^{18}x_{19} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 + 2^{16} \times 0 + 2^{17} \times 0 + 2^{18} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-1-4. 16 Variables, 2 Repeat Qubits, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]

For example of $n = 18$ [16 variables, and 2 repeat qubits], it is assumed that the logical formula, each value of $x_{1\sim 16}$, each value of $m_{1\sim 20}$, t , and k are same in the section 4-1-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000015259 (\approx 1/65536)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.0010767$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 159840, 254976, 163856, 241728, 32240, 257408, 0, 109568, 25104, 32856, 73744, 58448, 100356, 0, 17196, 54412, 0, 16640, 96104, 172160, 46860, 243744, 122848, 163968, 16448, 126720, 246912, 21456, 163140, 219416, 16512, 73720, 25088, 78592, 252800, 260248, 252864, 10208, 196608, 9320, 205568, 31232, 237584, 13332, 259456, 0, 4352, 260608, 82048, 108480$. (= spike)

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{18} = 262144$]] : $R_q \rightarrow$ candidates ; 159840 → 3, 5, 10, 13, 18, ... ; 254976 → 37, ... ; 163856 → 3, 5, 8, 16, ... ; 241728 → 13, 26, ... ; 32240 → 8, 16, ... ; 257408 → 55, ... ; 0 → nothingness ; 109568 → 2, 5, 7, 12, 24, ... ; 25104 → 10, 21, ... ; 32856 → 8, 16, ... ; 73744 → 4, 7, 14, 18, ... ; 58448 → 5, 9, 18, ... ; 100356 → 3, 5, 8, 13, 26, ... ; 17196 → 15, 30, ... ; 54412 → 5, 10, 14, 19, ... ; 16640 → 16, ... ; 96104 → 3, 6, 8, 11, 19, ... ; 172160 → 3, 6, 9, 12, 15, 17, ... ; 46860 → 6, 11, 17, ... ; 243744 → 14, 29, ... ; 122848 → 2, 4, 6, 9, 11, 13, 15, 17, ... ; 163968 → 3, 5, 8, 16, ... ; 16448 → 16, ... ; 126720 → 2, 4, 6, 8, 10, 12, 14, 17, ... ; 246912 → 17, ... ; 21456 → 12, 24, ... ; 163140 → 3, 5, 8, 16, ... ; 219416 → 6, 12, 18, ... ; 16512 → 16, ... ; 73720 → 4, 7, 14, 18, ... ; 25088 → 10, 21, ... ; 78592 → 3, 7, 10, 20, ... ; 252800 → 28, ... ; 260248 → 138, ... ; 252864 → 28, ... ; 10208 → 26, ... ; 196608 → 4, 8, 12, 16, ... ; 9320 → 28, ... ; 205568 → 5, 9, 14, 28, ... ; 31232 → 8, 17, ... ; 237584 → 11, 21, ... ; 13332 → 20, ... ; 259456 → 98, ... ; 4352 → 60, ... ; 260608 → 171, ... ; 82048 → 3, 6, 10, 13, 16, ... ; 108480 → 3, 5, 10, 12, 17,

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} + 2^{16}x_{17} + 2^{17}x_{18} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 + 2^{16} \times 0 + 2^{17} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-1-5. 16 Variables, 1 Repeat Qubit, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]

For example of $n = 17$ [16 variables, and 1 repeat qubit], it is assumed that the logical formula, each value of $x_{1\sim 16}$, each value of $m_{1\sim 20}$, t , and k are same in the section 4-1-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000015259 (\approx 1/65536)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.0012538$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 81984, 92166, 122848, 93656, 64, 5376, 125690, 118772, 123840, 119672, 121860, 0, 35936, 102368, 87936, 33814, 10272, 106112, 4096, 113120, 21510, 53200, 113712, 20480, 10110, 11898, 110486, 123390, 120792, 19460, 28234, 81624, 22144, 128514, 39416, 57472, 79200, 127152, 2048, 12802, 111040, 2050, 21304, 40912, 4592, 32256, 14784, 120816, 53356, 0. (= spike)$

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{17} = 131072$]] : $R_q \rightarrow$ candidates ; $81984 \rightarrow 3, 5, 8, 16, \dots$; $92166 \rightarrow 3, 7, 10, 17, \dots$; $122848 \rightarrow 16, \dots$; $93656 \rightarrow 4, 7, 14, 21, \dots$; $64 \rightarrow 2048, \dots$; $5376 \rightarrow 24, \dots$; $125690 \rightarrow 24, \dots$; $118772 \rightarrow 11, 21, \dots$; $123840 \rightarrow 18, \dots$; $119672 \rightarrow 12, 23, \dots$; $121860 \rightarrow 14, 28, \dots$; $0 \rightarrow$ nothingness ; $35936 \rightarrow 4, 7, 11, 22, \dots$; $102368 \rightarrow 5, 9, 18, \dots$; $87936 \rightarrow 3, 6, 9, 12, 15, 18, \dots$; $33814 \rightarrow 4, 8, 12, 16, \dots$; $10272 \rightarrow 13, 26, \dots$; $106112 \rightarrow 5, 11, 16, \dots$; $4096 \rightarrow 32, \dots$; $113120 \rightarrow 7, 15, 22, \dots$; $21510 \rightarrow 6, 12, 18, \dots$; $53200 \rightarrow 3, 5, 10, 15, 17, \dots$; $113712 \rightarrow 8, 15, 30, \dots$; $20480 \rightarrow 6, 13, 19, \dots$; $10110 \rightarrow 13, 26, \dots$; $11898 \rightarrow 11, 22, \dots$; $110486 \rightarrow 6, 13, 19, \dots$; $123390 \rightarrow 17, 34, \dots$; $120792 \rightarrow 13, 26, \dots$; $19460 \rightarrow 7, 13, 20, \dots$; $28234 \rightarrow 5, 9, 14, 28, \dots$; $81624 \rightarrow 3, 5, 8, 16, \dots$; $22144 \rightarrow 6, 12, 18, \dots$; $128514 \rightarrow 51, \dots$; $39416 \rightarrow 3, 7, 10, 20, \dots$; $57472 \rightarrow 2, 5, 7, 9, 16, \dots$; $79200 \rightarrow 3, 5, 10, 15, 20, \dots$; $127152 \rightarrow 33, \dots$; $2048 \rightarrow 64, \dots$; $12802 \rightarrow 10, 20, \dots$; $111040 \rightarrow 7, 13, 26, \dots$; $2050 \rightarrow 64, \dots$; $21304 \rightarrow 6, 12, 18, \dots$; $40912 \rightarrow 3, 6, 10, 13, 16, \dots$; $4592 \rightarrow 29, \dots$; $32256 \rightarrow 4, 8, 12, 16, \dots$; $14784 \rightarrow 9, 18, \dots$; $120816 \rightarrow 13, 26, \dots$; $53356 \rightarrow 3, 5, 10, 15, 17, \dots$.

When u is $16 (2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} + 2^{16}x_{17} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 + 2^{16} \times 0 = 16)$, the value of logical formula is 1. Therefore, it is the answer.

4-1-6. 16 Variables, 0 Repeat Qubit, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]

For example of $n = 16$ [16 variables, and 0 repeat qubit], it is assumed that the logical formula, each value of $x_{1\sim 16}$, each value of $m_{1\sim 20}$, t , and k are same in the section 4-1-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000015259 (\approx 1/65536)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.0012525$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 2156, 12256, 60671, 2596, 8964, 61968, 416, 4032, 59393, 61439, 57285, 41984, 6145, 1280, 51396, 63454,$

41264, 55616, 28096, 34560, 11726, 61706, 12916, 4484, 63728, 0, 52144, 0, 5108, 8287, 59418, 64508, 65264, 2047, 60984, 22272, 44961, 8312, 62270, 1024, 4058, 0, 4416, 26328, 65525, 62017, 8138, 56328, 53756, 63638. (= spike)

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{16} = 65536$]] : $R_q \rightarrow$ candidates ; 2156 \rightarrow 30, ... ; 12256 \rightarrow 5, 11, 16, ... ; 60671 \rightarrow 13, 27, ... ; 2596 \rightarrow 25, ... ; 8964 \rightarrow 7, 15, 22, ... ; 61968 \rightarrow 18, 37, ... ; 416 \rightarrow 158, ... ; 4032 \rightarrow 16, ... ; 59393 \rightarrow 11, 21, ... ; 61439 \rightarrow 16, ... ; 57285 \rightarrow 8, 16, ... ; 41984 \rightarrow 3, 6, 8, 11, 14, 25, ... ; 6145 \rightarrow 11, 21, ... ; 1280 \rightarrow 51, ... ; 51396 \rightarrow 5, 9, 14, 28, ... ; 63454 \rightarrow 31, ... ; 41264 \rightarrow 3, 5, 8, 16, ... ; 55616 \rightarrow 7, 13, 20, ... ; 28096 \rightarrow 2, 5, 7, 14, 21, ... ; 34560 \rightarrow 2, 4, 6, 8, 11, 13, 15, 17, ... ; 11726 \rightarrow 6, 11, 17, ... ; 61706 \rightarrow 17, ... ; 12916 \rightarrow 5, 10, 15, 20, ... ; 4484 \rightarrow 15, 29, ... ; 63728 \rightarrow 36, ... ; 0 \rightarrow nothingness ; 52144 \rightarrow 5, 10, 15, 20, ... ; 5108 \rightarrow 13, 26, ... ; 8287 \rightarrow 8, 16, ... ; 59418 \rightarrow 11, 21, ... ; 64508 \rightarrow 64, ... ; 65264 \rightarrow 241, ... ; 2047 \rightarrow 32, ... ; 60984 \rightarrow 14, 29, ... ; 22272 \rightarrow 3, 6, 9, 12, 15, 18, ... ; 4496 \rightarrow 3, 6, 10, 13, 16, ... ; 8312 \rightarrow 8, 16, ... ; 62270 \rightarrow 20, ... ; 1024 \rightarrow 64, ... ; 4058 \rightarrow 16, ... ; 4416 \rightarrow 15, 30, ... ; 26328 \rightarrow 3, 5, 10, 15, 20, ... ; 65525 \rightarrow 5958, ... ; 62017 \rightarrow 19, ... ; 8138 \rightarrow 8, 16, ... ; 56328 \rightarrow 7, 14, 21, ... ; 53756 \rightarrow 6, 11, 22, ... ; 63638 \rightarrow 35,

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-2. 15 Variables

4-2-1. 15 Variables, 6 Repeat Qubits, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]

For example of $n = 21$ [15 variables, and 6 repeat qubits], it is assumed that a logical formula : $(x_3 \vee x_4 \vee x_5) \& (\sim x_1 \vee x_2 \vee x_3) \& (\sim x_3 \vee x_4 \vee x_5) \& (x_3 \vee \sim x_4 \vee x_5) \& (\sim x_2 \vee x_3 \vee \sim x_5) \& (\sim x_3 \vee \sim x_4 \vee x_5) \& (\sim x_3 \vee x_4 \vee \sim x_5) \& (x_3 \vee \sim x_4 \vee \sim x_5) \& (\sim x_3 \vee \sim x_4 \vee \sim x_5) \& (x_4 \vee \sim x_5 \vee \sim x_6) \& (\sim x_5 \vee x_6 \vee \sim x_7) \& (x_6 \vee x_7 \vee \sim x_8) \& (x_7 \vee x_8 \vee \sim x_9) \& (x_8 \vee x_9 \vee \sim x_{10}) \& (x_9 \vee x_{10} \vee \sim x_{11}) \& (x_{10} \vee x_{11} \vee \sim x_{12}) \& (x_{11} \vee x_{12} \vee \sim x_{13}) \& (x_{12} \vee x_{13} \vee \sim x_{14}) \& (x_{13} \vee x_{14} \vee \sim x_{15})$, each value of x_{1-15} : $x_1 = x_2 = x_3 = x_4 = x_6 = x_7 = x_8 = x_9 = x_{10} = x_{11} = x_{12} = x_{13} = x_{14} = x_{15} = 0$, $x_5 = 1$, $m_1 = 1$, $m_2 = 2$, $m_3 = 3$, $m_4 = 4$, $m_5 = 5$, $m_6 = 6$, $m_7 = 7$, $m_8 = 8$, $m_9 = 9$, $m_{10} = 10$, $m_{11} = 11$, $m_{12} = 12$, $m_{13} = 13$, $m_{14} = 14$, $m_{15} = 15$, $m_{16} = 16$, $m_{17} = 17$, $m_{18} = 18$, $m_{19} = 19$, $t = 7$, and $k = (19 + 1)19/2 = 190$.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000030518 (\approx 1/32768)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.000000000$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 1306624, 1064960, 405376, 346624, 1364160, 2054080, 458240, 656128, 2097024, 2078080, 295168, 1998976, 1011712, 386560, 2096640, 269952, 175168, 1647104, 1840960, 768512, 0, 556672, 35328, 162816, 1840704, 2046208, 20096, 1912832, 280576, 2031872, 1220608, 16384, 441856, 110976, 1224768, 0, 1040320, 456704, 330752, 2035712, 1769408, 2090496, 182272, 1837248, 1247744, 192384, 1469056, 917504, 1898688, 57344. (= spike)$

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{21} = 2097152$]] : $R_q \rightarrow$ candidates ; 1306624 \rightarrow 3, 5, 8, 16, ... ; 1064960 \rightarrow 2, 4, 6, 8, 10, 12, 14, 16, ... ; 405376 \rightarrow 5, 10, 16, ... ; 346624 \rightarrow 6, 12, 18, ... ; 1364160 \rightarrow 3, 6, 9, 11, 14, 17, ... ; 2054080 \rightarrow 49, ... ; 458240 \rightarrow 5, 9, 18, ... ; 656128 \rightarrow 3, 6, 10, 13, 16, ... ; 2097024 \rightarrow 16384, ... ; 2078080 \rightarrow 110, ... ; 295168 \rightarrow 7, 14, 21, ... ; 1998976 \rightarrow 21, ... ; 1011712 \rightarrow 2, 4, 6, 8, 10, 12, 15, 17, ... ; 386560 \rightarrow 5, 11, 22, ... ; 2096640 \rightarrow 4096, ... ; 269952 \rightarrow 8, 16, ... ; 175168 \rightarrow 12, 24, ... ; 1647104 \rightarrow 5, 9, 14, 28, ... ; 1840960 \rightarrow 8, 16, ... ; 768512 \rightarrow 3, 6, 8, 11, 19, ... ; 0 \rightarrow nothingness ; 556672 \rightarrow 4, 8, 11, 15, 30, ... ; 35328 \rightarrow 59, ... ; 162816 \rightarrow 13, 26, ... ; 1840704 \rightarrow 8, 16, ... ; 2046208 \rightarrow 41, ... ; 20096 \rightarrow 104, ... ; 1912832 \rightarrow 11, 23, ... ; 280576 \rightarrow 8, 15, 30, ... ; 2031872 \rightarrow 32, ... ; 1220608 \rightarrow 2, 5, 7, 12, 24, ... ; 16384 \rightarrow 128, ... ; 441856 \rightarrow 5, 10, 14, 19, ... ; 110976 \rightarrow 19, ... ; 1224768 \rightarrow 3, 5, 7, 12, 24, ... ; 1040320 \rightarrow 2, 4, 6, 8, 10, 12, 14, 16, ... ; 456704 \rightarrow 5, 9, 14, 23, ... ; 330752 \rightarrow 6, 13, 19, ... ; 2035712 \rightarrow 34, ... ; 1769408 \rightarrow 6, 13, 19, ... ; 2090496 \rightarrow 315, ... ; 182272 \rightarrow 12, 23, ... ; 1837248 \rightarrow 8, 16, ... ; 1247744 \rightarrow 3, 5, 10, 15, 20, ... ; 192384 \rightarrow 11, 22, ... ; 1469056 \rightarrow 3, 7, 10, 20, ... ; 917504 \rightarrow 2, 5, 7, 9, 16, ... ; 1898688 \rightarrow 11, 21, ... ; 57344 \rightarrow 37,

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} + 2^{16}x_{17} + 2^{17}x_{18} + 2^{18}x_{19} + 2^{19}x_{20} + 2^{20}x_{21} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 + 2^{16} \times 0 + 2^{17} \times 0 + 2^{18} \times 0 + 2^{19} \times 0 + 2^{20} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-2-2. 15 Variables, 5 Repeat Qubits, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19]

For example of $n = 20$ [15 variables, and 5 repeat qubits], it is assumed that the logical formula, each value of $x_{1\sim 15}$, each value of $m_{1\sim 19}$, t , and k are same in the section 4-2-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000030518 (\approx 1/32768)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.000000000$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 458496, 488448, 593216, 985600, 987328, 310272, 352256, 17152, 933888, 1033408, 1001984, 370688, 33280, 850688, 721024, 1024000, 28416, 704000, 196928, 620544, 0, 646656, 1015296, 41984, 802560, 941824, 40864, 688288, 16896, 126976, 933888, 1015744, 61376, 868352, 114816, 659456, 966752, 33792, 16128, 656128, 34048, 982784, 425728, 962560, 792128, 900864, 20352, 492704, 73472, 131168. (= spike)$

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{20} = 1048576$]] : $R_q \rightarrow$ candidates ; 458496 \rightarrow 2, 5, 7, 9, 16, ... ; 488448 \rightarrow 2, 4, 6, 9, 11, 13, 15, 30, ... ; 593216 \rightarrow 2, 5, 7, 14, 16, ... ; 985600 \rightarrow 17, ... ; 987328 \rightarrow 17, ... ; 310272 \rightarrow 3, 7, 10, 17, ... ; 352256 \rightarrow 3, 6, 9, 12, 15, 18, ... ; 17152 \rightarrow 61, ... ; 933888 \rightarrow 9, 18, ... ; 1033408 \rightarrow 69, ... ; 1001784 \rightarrow 22, ... ; 370688 \rightarrow 3, 6, 9, 11, 14, 17, ... ; 33280 \rightarrow 32, ... ; 850688 \rightarrow 5, 11, 16, ... ; 721024 \rightarrow 3, 6, 10, 13, 16, ... ; 1024000

→ 43, ... ; 28416 → 37, ... ; 704000 → 3, 6, 9, 12, 15, 18, ... ; 196928 → 5, 11, 16, ... ; 620544 → 3, 5, 10, 15, 17, ... ; 0 → nothingness ; 646656 → 3, 5, 8, 13, 26, ... ; 1015296 → 32, ... ; 41984 → 25, ... ; 802560 → 4, 9, 13, 17, ... ; 941824 → 10, 20, ... ; 40864 → 26, ... ; 688288 → 3, 6, 9, 12, 15, 17, ... ; 16896 → 62, ... ; 126976 → 8, 17, ... ; 933888 → 9, 18, ... ; 1015744 → 32, ... ; 61376 → 17, ... ; 868352 → 6, 12, 17, ... ; 114816 → 9, 18, ... ; 659456 → 3, 5, 8, 16, ... ; 966752 → 13, 26, ... ; 33792 → 31, ... ; 16128 → 65, ... ; 656128 → 3, 5, 8, 16, ... ; 34048 → 31, ... ; 982784 → 16, ... ; 425728 → 3, 5, 10, 15, 17, ... ; 962560 → 12, 24, ... ; 792128 → 4, 8, 12, 16, ... ; 900864 → 7, 14, 21, ... ; 20352 → 52, ... ; 492704 → 2, 4, 7, 9, 11, 20, ... ; 73472 → 14, 29, ... ; 131168 → 8, 16,

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} + 2^{16}x_{17} + 2^{17}x_{18} + 2^{18}x_{19} + 2^{19}x_{20} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 + 2^{16} \times 0 + 2^{17} \times 0 + 2^{18} \times 0 + 2^{19} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-2-3. 15 Variables, 4 Repeat Qubits, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]

For example of $n = 19$ [15 variables, and 4 repeat qubits], it is assumed that the logical formula, each value of x_{1-15} , each value of m_{1-19} , t , and k are same in the section 4-2-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000030518 (\approx 1/32768)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.00093793$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 274848, 167680, 508160, 487424, 12800, 146944, 40928, 505920, 522944, 24576, 16192, 523776, 475904, 356352, 523264, 310272, 184576, 188544, 16912, 163616, 490416, 520512, 832, 824000, 68736, 32960, 510464, 479232, 19456, 16480, 523248, 24528, 5120, 38272, 371120, 360032, 421792, 16256, 169728, 32752, 27136, 506560, 507904, 34816, 31232, 490496, 503552, 507968, 360704, 524032. (= spike)$

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{19} = 524288$]] : $R_q \rightarrow$ candidates ; 274848 → 2, 4, 6, 8, 11, 13, 15, 17, ... ; 167680 → 3, 6, 9, 13, 16, ... ; 508160 → 33, ... ; 487424 → 14, 28, ... ; 12800 → 41, ... ; 146944 → 4, 7, 14, 18, ... ; 40928 → 13, 26, ... ; 505920 → 29, ... ; 522944 → 390, ... ; 24576 → 21, ... ; 16192 → 32, ... ; 523776 → 1024, ... ; 475904 → 11, 22, ... ; 356352 → 3, 6, 9, 13, 16, ... ; 523264 → 512, ... ; 310272 → 3, 5, 10, 15, 17, ... ; 184576 → 3, 6, 9, 11, 14, 17, ... ; 188544 → 3, 6, 8, 11, 14, 25, ... ; 16912 → 31, ... ; 163616 → 3, 6, 10, 13, 16, ... ; 490416 → 15, 31, ... ; 520512 → 139, ... ; 832 → 630, ... ; 82400 → 6, 13, 19, ... ; 68736 → 8, 15, 23, ... ; 32960 → 16, ... ; 510464 → 38, ... ; 479232 → 12, 23, ... ; 19456 → 27, ... ; 16480 → 32, ... ; 523248 → 504, ... ; 24528 → 21, ... ; 5120 → 102, ... ; 38272 → 14, 27, ... ; 371120 → 3, 7, 14, 17, ... ; 360032 → 3, 6, 10, 13, 16, ... ; 421792 → 5, 10, 15, 20, ... ; 16256 → 32, ... ; 169728 → 3, 6, 9, 12, 15, 19, ... ; 32752 → 16, ... ; 27136 → 19, ... ; 506560 → 30, ... ; 507904 → 32, ...

; 34816 \rightarrow 15, 30, ... ; 31232 \rightarrow 17, ... ; 490496 \rightarrow 16, ... ; 503552 \rightarrow 25, ... ; 507968 \rightarrow 32, ... ; 360704 \rightarrow 3, 6, 10, 13, 16, ... ; 524032 \rightarrow 2048,

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} + 2^{16}x_{17} + 2^{17}x_{18} + 2^{18}x_{19} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 + 2^{16} \times 0 + 2^{17} \times 0 + 2^{18} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-2-4. 15 Variables, 3 Repeat Qubits, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]

For example of $n = 18$ [15 variables, and 3 repeat qubits], it is assumed that the logical formula, each value of $x_{1\sim 15}$, each value of $m_{1\sim 19}$, t , and k are same in the section 4-2-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000030518 (\approx 1/32768)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.00053154$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 12352, 121344, 9216, 245120, 168960, 67584, 16632, 254176, 640, 43008, 137320, 8752, 23040, 241152, 13296, 250784, 246784, 164672, 32704, 69632, 253568, 171776, 32256, 188736, 108480, 261440, 71392, 45056, 59392, 214272, 254464, 258848, 26240, 222216, 10048, 203552, 49144, 106752, 254016, 180352, 78632, 16608, 254688, 46056, 48288, 254208, 246016, 229912, 12288, 257664$. (= spike)

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{18} = 262144$]] : $R_q \rightarrow$ candidates ; 12352 \rightarrow 21, ... ; 121344 \rightarrow 2, 4, 7, 9, 11, 13, 26, ... ; 9216 \rightarrow 28, ... ; 245120 \rightarrow 15, 31, ... ; 168960 \rightarrow 3, 6, 8, 11, 14, 28, ... ; 67584 \rightarrow 4, 8, 12, 16, ... ; 254176 \rightarrow 33, ... ; 640 \rightarrow 410, ... ; 43008 \rightarrow 6, 12, 18, ... ; 137320 \rightarrow 2, 4, 6, 8, 11, 13, 15, 17, ... ; 8752 \rightarrow 30, ... ; 23040 \rightarrow 11, 23, ... ; 241152 \rightarrow 13, 25, ... ; 13296 \rightarrow 20, ... ; 250784 \rightarrow 23, ... ; 246784 \rightarrow 17, ... ; 164672 \rightarrow 3, 5, 8, 16, ... ; 32704 \rightarrow 8, 16, ... ; 69632 \rightarrow 4, 8, 11, 15, 30, ... ; 253568 \rightarrow 31, ... ; 171776 \rightarrow 3, 6, 9, 12, 15, 17, ... ; 32256 \rightarrow 8, 16, ... ; 188736 \rightarrow 4, 7, 14, 18, ... ; 108480 \rightarrow 3, 5, 10, 12, 17, ... ; 261440 \rightarrow 372, ... ; 71392 \rightarrow 4, 7, 11, 22, ... ; 45056 \rightarrow 6, 12, 17, ... ; 59392 \rightarrow 4, 9, 13, 22, ... ; 214272 \rightarrow 6, 11, 22, ... ; 254464 \rightarrow 34, ... ; 258848 \rightarrow 80, ... ; 26240 \rightarrow 10, 20, ... ; 222216 \rightarrow 7, 13, 26, ... ; 10048 \rightarrow 26, ... ; 203552 \rightarrow 5, 9, 18, ... ; 49144 \rightarrow 5, 11, 16, ... ; 106752 \rightarrow 3, 5, 10, 15, 17, ... ; 254016 \rightarrow 32, ... ; 180352 \rightarrow 3, 6, 10, 13, 16, ... ; 78632 \rightarrow 3, 7, 10, 20, ... ; 16608 \rightarrow 16, ... ; 254688 \rightarrow 35, ... ; 46056 \rightarrow 6, 11, 17, ... ; 48288 \rightarrow 5, 11, 22, ... ; 254208 \rightarrow 33, ... ; 246016 \rightarrow 16, ... ; 229912 \rightarrow 8, 16, ... ; 12288 \rightarrow 21, ... ; 257664 \rightarrow 59,

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} + 2^{16}x_{17} + 2^{17}x_{18} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 + 2^{16} \times 0 + 2^{17} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-2-5. 15 Variables, 2 Repeat Qubits, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]

For example of $n = 17$ [15 variables, and 2 repeat qubits], it is assumed that the logical formula, each value of $x_{1\sim 15}$, each value of $m_{1\sim 19}$, t , and k are same in the section 4-2-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000030518 (\approx 1/32768)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.0013206$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 114732, 95216, 20412, 72320, 15552, 90104, 47104, 131040, 122624, 70656, 7424, 106464, 3752, 106776, 0, 5952, 122016, 123904, 3080, 130048, 98464, 130848, 114752, 113824, 12208, 2240, 116732, 40928, 127424, 122720, 114016, 85848, 124768, 123136, 8120, 65792, 84952, 125776, 130048, 116736, 47136, 126576, 4128, 116236, 114544, 107264, 17860, 63296, 85888, 122560$. (= spike)

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{17} = 131072$]] : $R_q \rightarrow$ candidates ; $114732 \rightarrow 8, 16, \dots$; $95216 \rightarrow 4, 7, 11, 22, \dots$; $20412 \rightarrow 6, 13, 26, \dots$; $72320 \rightarrow 2, 5, 7, 9, 18, \dots$; $15552 \rightarrow 8, 17, \dots$; $90104 \rightarrow 3, 6, 10, 13, 16, \dots$; $47104 \rightarrow 3, 6, 8, 11, 14, 25, \dots$; $131040 \rightarrow 4096, \dots$; $122624 \rightarrow 16, \dots$; $70656 \rightarrow 2, 4, 7, 9, 11, 13, 26, \dots$; $7424 \rightarrow 18, \dots$; $106464 \rightarrow 5, 11, 16, \dots$; $3752 \rightarrow 35, \dots$; $106776 \rightarrow 5, 11, 16, \dots$; $0 \rightarrow$ nothingness ; $5952 \rightarrow 22, \dots$; $122016 \rightarrow 14, 29, \dots$; $123904 \rightarrow 18, \dots$; $3080 \rightarrow 43, \dots$; $130048 \rightarrow 128, \dots$; $98464 \rightarrow 4, 8, 12, 16, \dots$; $130848 \rightarrow 585, \dots$; $114752 \rightarrow 8, 16, \dots$; $113824 \rightarrow 8, 15, 23, \dots$; $12208 \rightarrow 11, 21, \dots$; $2240 \rightarrow 59, \dots$; $116732 \rightarrow 9, 18, \dots$; $40928 \rightarrow 3, 6, 10, 13, 16, \dots$; $127424 \rightarrow 36, \dots$; $122720 \rightarrow 16, \dots$; $114016 \rightarrow 8, 15, 23, \dots$; $85848 \rightarrow 3, 6, 9, 12, 15, 17, \dots$; $124768 \rightarrow 21, \dots$; $123136 \rightarrow 17, \dots$; $8120 \rightarrow 16, \dots$; $65792 \rightarrow 2, 4, 6, 8, 10, 12, 14, 16, \dots$; $84952 \rightarrow 3, 6, 9, 11, 14, 17, \dots$; $125776 \rightarrow 25, \dots$; $130048 \rightarrow 128, \dots$; $116736 \rightarrow 9, 18, \dots$; $47136 \rightarrow 3, 6, 8, 11, 14, 25, \dots$; $126576 \rightarrow 29, \dots$; $4128 \rightarrow 32, \dots$; $116236 \rightarrow 9, 18, \dots$; $114544 \rightarrow 8, 16, \dots$; $107264 \rightarrow 6, 11, 22, \dots$; $17860 \rightarrow 7, 15, 22, \dots$; $63296 \rightarrow 2, 4, 6, 8, 10, 12, 15, 17, \dots$; $85888 \rightarrow 3, 6, 9, 12, 15, 17, \dots$; $122560 \rightarrow 15, 31, \dots$.

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} + 2^{16}x_{17} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 + 2^{16} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-2-6. 15 Variables, 1 Repeat Qubit, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]

For example of $n = 16$ [15 variables, and 1 repeat qubit], it is assumed that the logical formula, each value of $x_{1\sim 15}$, each value of $m_{1\sim 19}$, t , and k are same in the section 4-2-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000030518 (\approx 1/32768)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.0014779$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 20192, 61952, 1536, 4230, 16288, 64464, 44976, 62016, 28632, 64896, 10456, 8076, 6080, 56316, 65280, 55302, 62720, 39190, 104, 35636, 47936, 63104, 5632, 5024, 61240, 11520, 63616, 12904, 0, 6, 20464, 11244, 43584, 58720, 47618, 4096, 62224, 8340, 4084, 3520, 528, 20992, 2044, 976, 0, 60416, 54272, 64000, 58112, 0.$ (= spike)

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{16} = 65536$]] : $R_q \rightarrow$ candidates ; $20192 \rightarrow 3, 7, 10, 13, 26, \dots$; $61952 \rightarrow 18, \dots$; $1536 \rightarrow 21, \dots$; $4230 \rightarrow 8, 16, \dots$; $16288 \rightarrow 2, 4, 6, 8, 10, 12, 14, 16, \dots$; $64464 \rightarrow 31, \dots$; $44976 \rightarrow 3, 5, 8, 16, \dots$; $62016 \rightarrow 9, 19, \dots$; $28632 \rightarrow 8, 16, \dots$; $64896 \rightarrow 51, \dots$; $10456 \rightarrow 3, 6, 9, 13, 16, \dots$; $8076 \rightarrow 4, 8, 12, 16, \dots$; $6080 \rightarrow 5, 11, 16, \dots$; $56316 \rightarrow 4, 7, 14, 18, \dots$; $65280 \rightarrow 128, \dots$; $55302 \rightarrow 3, 6, 10, 13, 16, \dots$; $62720 \rightarrow 12, 23, \dots$; $39190 \rightarrow 5, 10, 15, 20, \dots$; $104 \rightarrow 315, \dots$; $35636 \rightarrow 11, 23, \dots$; $47936 \rightarrow 2, 4, 7, 9, 11, 13, 26, \dots$; $63104 \rightarrow 13, 27, \dots$; $5632 \rightarrow 6, 12, 17, \dots$; $5024 \rightarrow 7, 13, 26, \dots$; $61240 \rightarrow 8, 15, 23, \dots$; $11520 \rightarrow 3, 6, 9, 11, 14, 17, \dots$; $63616 \rightarrow 17, \dots$; $12904 \rightarrow 3, 5, 10, 15, 18, \dots$; $0 \rightarrow$ nothingness ; $6 \rightarrow 5461, \dots$; $20464 \rightarrow 3, 5, 8, 16, \dots$; $11244 \rightarrow 3, 6, 9, 12, 15, 17, \dots$; $43584 \rightarrow 3, 6, 9, 12, 15, 18, \dots$; $58720 \rightarrow 5, 10, 14, 19, \dots$; $47618 \rightarrow 2, 4, 7, 9, 11, 22, \dots$; $4096 \rightarrow 8, 16, \dots$; $62224 \rightarrow 10, 20, \dots$; $8340 \rightarrow 4, 8, 12, 16, \dots$; $4084 \rightarrow 8, 16, \dots$; $3520 \rightarrow 9, 19, \dots$; $528 \rightarrow 62, \dots$; $20992 \rightarrow 3, 6, 8, 11, 14, 25, \dots$; $2044 \rightarrow 16, \dots$; $976 \rightarrow 34, \dots$; $60416 \rightarrow 6, 13, 19, \dots$; $54272 \rightarrow 3, 6, 9, 12, 15, 17, \dots$; $64000 \rightarrow 21, \dots$; $58112 \rightarrow 4, 9, 13, 22, \dots$.

When u is 16 ($2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} + 2^{15}x_{16} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 + 2^{15} \times 0 = 16$), the value of logical formula is 1. Therefore, it is the answer.

4-2-7. 15 Variables, 0 Repeat Qubit, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]

For example of $n = 15$ [15 variables, and 0 repeat qubit], it is assumed that the logical formula, each value of x_{1-15} , each value of m_{1-19} , t , and k are same in the section 4-2-1.

A result of this problem is the following.

The probability probe value of $|w_{1,j}\rangle = 0 : \approx 0.000030518 (\approx 1/32768)$.

The probability probe value of $|x_i\rangle = 16 : \approx 0.0014756$.

The example of 50 times test : The read value of $|x_i\rangle$; $R_q = 6152, 19532, 31092, 32064, 11392, 26590, 23040, 3064, 2080, 31616, 32741, 2071, 4096, 29376, 12478, 4104, 926, 28672, 25976, 2260, 3968, 19481, 32258, 28580, 5600, 4064, 27120, 4474, 22400, 3440, 21552, 1804, 32256, 4600, 136, 1248, 27632, 31328, 1409, 18488, 31512, 22652, 31028, 27360, 30832, 0, 21936, 21308, 48, 29824.$ (= spike)

The candidates of number of spikes are estimated by the function [the function estimate_num_spikes (spike, range) [spike : read value, range : $2^n = 2^{15} = 32768$]] : $R_q \rightarrow$ candidates ; $6152 \rightarrow 5, 11, 16, \dots$; $19532 \rightarrow 3, 5, 10, 15, 20, \dots$; $31092 \rightarrow 20, \dots$; $32064 \rightarrow 47, \dots$; $11392 \rightarrow 3, 6, 9, 12, 14, 17, \dots$; $26590 \rightarrow 5, 11, 16, \dots$; $23040 \rightarrow 3, 7, 10, 17, \dots$; $3064 \rightarrow 11, 21, \dots$; $2080 \rightarrow 16, \dots$; $31616 \rightarrow 28, \dots$; $32741 \rightarrow 1214, \dots$; $2072 \rightarrow 16, \dots$; $4096 \rightarrow 8, 16, \dots$; $29376 \rightarrow 10, 19, \dots$; $12478 \rightarrow 3, 5, 8,$

13, 21, ... ; 4104 \rightarrow 8, 16, ... ; 926 \rightarrow 35, ... ; 28672 \rightarrow 8, 16, ... ; 25976 \rightarrow 5, 10, 14, 19, ... ; 2260 \rightarrow 15, 29, ... ; 3968 \rightarrow 8, 17, ... ; 19481 \rightarrow 3, 5, 10, 15, 20, ... ; 32258 \rightarrow 64, ... ; 28580 \rightarrow 8, 16, ... ; 5600 \rightarrow 6, 12, 18, ... ; 4064 \rightarrow 8, 16, ... ; 27120 \rightarrow 6, 12, 17, ... ; 4474 \rightarrow 7, 15, 22, ... ; 22400 \rightarrow 3, 6, 10, 13, 16, ... ; 3440 \rightarrow 10, 19, ... ; 21552 \rightarrow 3, 6, 9, 12, 15, 18, ... ; 1804 \rightarrow 18, ... ; 32256 \rightarrow 64, ... ; 4600 \rightarrow 7, 14, 21, ... ; 136 \rightarrow 241, ... ; 1248 \rightarrow 26, ... ; 27632 \rightarrow 6, 13, 19, ... ; 31328 \rightarrow 23, ... ; 1409 \rightarrow 23, ... ; 18488 \rightarrow 2, 5, 7, 14, 16, ... ; 31512 \rightarrow 216, ... ; 22652 \rightarrow 3, 7, 10, 13, 26, ... ; 31028 \rightarrow 19, ... ; 27360 \rightarrow 6, 12, 18, ... ; 30832 \rightarrow 17, ... ; 0 \rightarrow nothingness ; 21936 \rightarrow 3, 6, 9, 12, 15, 18, ... ; 21308 \rightarrow 3, 6, 9, 11, 14, 17, ... ; 48 \rightarrow 683, ... ; 29824 \rightarrow 11, 22,

When u is $16(2^0x_1 + 2^1x_2 + 2^2x_3 + 2^3x_4 + 2^4x_5 + 2^5x_6 + 2^6x_7 + 2^7x_8 + 2^8x_9 + 2^9x_{10} + 2^{10}x_{11} + 2^{11}x_{12} + 2^{12}x_{13} + 2^{13}x_{14} + 2^{14}x_{15} = 2^0 \times 0 + 2^1 \times 0 + 2^2 \times 0 + 2^3 \times 0 + 2^4 \times 1 + 2^5 \times 0 + 2^6 \times 0 + 2^7 \times 0 + 2^8 \times 0 + 2^9 \times 0 + 2^{10} \times 0 + 2^{11} \times 0 + 2^{12} \times 0 + 2^{13} \times 0 + 2^{14} \times 0 = 16)$, the value of logical formula is 1. Therefore, it is the answer.

5. Discussion

In the 3-SAT problem, when [(the logical formula) = 1] is obtained, there is only one combination.

5-1. 16 Variables

5-1-1. 16 Variables, Repeat Qubits from 0 to 5, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]

In the section 4-1, there is RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]. And then, [(the logical formula) = 1] of combination of variables is selected, and the computation of repeats is done. When N is 2^{16} , in the Grover's method, the complexity is $N^{1/2} = 2^{16/2} = 256$, in the new method, for (variables, repeat qubits) = (16, 0), it is $50/10 = 5$, for (16, 1), it is $50/8 \approx 6$, for (16, 2), it is $50/10 = 5$, for (16, 3), it is $50/8 \approx 6$, for (16, 4), it is $50/6 \approx 8$, and for (16, 5), it is $50/5 = 10$.

In this range, the new method is less than the complexity of the Grover's method, and then, in for (16, 0), and (16, 2), the probability is the maximum value 20%.

5-1-2. 16 Variables, Repeat Qubits from 0 to 5, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 1]

There are RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 1], $t = 7$, and $k = 191$. And then, [(the logical formula) = 1] of combination of variables is selected, and the computation of repeats is done. In the new method, for (16, 0), it is $50/6 \approx 8$, for (16, 1), it is $50/8 \approx 6$, for (16, 2), it is $50/10 = 5$, for (16, 3), it is $50/6 \approx 8$, for (16, 4), it is $50/10 = 5$, and for (16, 5), it is $50/8 \approx 6$.

In this range, the new method is less than the complexity of the Grover's method, and then, in for (16, 2), and (16, 4), the probability is the maximum value 20%.

5-1-3. 16 Variables, Repeat Qubits from 0 to 5, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 1, 2]

There are RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 1, 2], $t = 7$, and $k = 174$. And then, [(the logical formula) = 1] of combination of variables is selected, and the computation of repeats is done. In the new method, for (16, 0), it is

$50/11 \approx 5$, for (16, 1), it is $50/10 = 5$, for (16, 2), it is $50/8 \approx 6$, for (16, 3), it is $50/11 \approx 5$, for (16, 4), it is $50/4 = 13$, and for (16, 5), it is $50/10 = 5$.

In this range, the new method is less than the complexity of the Grover's method, and then, in for (16, 0), and (16, 3), the probability is the maximum value 22%.

5-1-4. 16 Variables, Repeat Qubits from 0 to 5, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 1, 2, 3]

There are RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 1, 2, 3], $t = 7$, and $k = 159$. And then, [(the logical formula) = 1] of combination of variables is selected, and the computation of repeats is done. In the new method, for (16, 0), it is $50/12 \approx 4$, for (16, 1), it is $50/13 \approx 4$, for (16, 2), it is $50/9 \approx 6$, for (16, 3), it is $50/13 \approx 4$, for (16, 4), it is $50/6 \approx 8$, and for (16, 5), it is $50/6 \approx 8$.

In this range, the new method is less than the complexity of the Grover's method, and then, in for (16, 1), and (16, 3), the probability is the maximum value 26%.

From step 11 of section 3 (Quantum Algorithm), the next weight changed computation isn't done.

5-2. 15Variables

5-2-1. 15 Variables, Repeat Qubits from 0 to 6, and RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]

In the section 4-2, there is RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]. And then, [(the logical formula) = 1] of combination of variables is selected, and the computation of repeats is done. When N is 2^{15} , in the Grover's method, the complexity is $N^{1/2} = 2^{15/2} \approx 181$, in the new method, for (15, 0), it is $50/11 \approx 5$, for (15, 1), it is $50/13 \approx 4$, for (15, 2), it is $50/12 \approx 4$, for (15, 3), it is $50/10 = 5$, for (15, 4), it is $50/8 \approx 6$, for (15, 5), it is $50/10 = 5$, and for (15, 6), it is $50/10 = 5$.

In this range, the new method is less than the complexity of the Grover's method, and then, in for (15, 1), the probability is the maximum value 26%.

From step 11 of section 3 (Quantum Algorithm), the next weight changed computation isn't done.

Summary

The quantum algorithm for the 3-SAT problem of 15, and 16 variables by the quantum Fourier transform with the repeat qubits, and the weight changes on the QC Engine, and its example are reported.

The complexity of this method is several times, and then, in for (15, 1) of RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19], the probability is the maximum value 26%, and in for (16, 1), and (16, 3) of RAM = [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 1, 2, 3], the probability is the maximum value 26%.

I will apply this method for other problems.

References

[1] Deutsch, D., and Jozsa, R., 1992, "Rapid solution of problems by quantum computation," Proc. Roy. Soc. Lond. A **439**, 553-558.

- [2] Shor, P. W., 1994, "Algorithms for quantum computation : discrete logarithms and factoring," Proc. 35th Annu. Symp. Foundations of Computer Science, IEEE, pp.124-134.
- [3] Grover, L. K., 1996, "A fast quantum mechanical algorithm for database search," Proc. 28th Annu. ACM Symp. Theory of Computing, pp.212-219.
- [4] Grover, L. K., 1998, "A framework for fast quantum mechanical algorithms," Proc. 30th Annu. ACM Symp. Theory of Computing, pp.53-62.
- [5] Cook, S. A., 1971, "The complexity of theorem proving procedures," Proc. 3rd Annu. ACM Symp. Theory of Computing, pp.151-158.
- [6] Johnston, E.R., Harrigan, N., and Gimeno-Segovia, M., 2019, Programming Quantum Computers, O'Reilly, ISBN 978-1-492-03968-6.
- [7] Fujimura, T., 2024, "Quantum algorithm for 3-SAT problem by Shor's Fourier transform with RAM on QCEngine," Glob. J. Pure Appl. Math., **20**, 695-703.
- [8] Fujimura, T., 2026, "Quantum algorithm for 3-SAT problem of 14 variables by quantum Fourier transform with repeat qubits, and weight changes on QCEngine," Glob. J. Pure Appl. Math., **22**, 85-99.
- [9] Takeuchi, S., 2005, Ryoshi Konpyuta (Quantum Computer), Kodansha, Tokyo, Japan [in Japanese].
- [10] Miyano, K., and Furusawa, A., 2008, Ryoshi Konpyuta Nyumon (An Introduction to Quantum Computation), Nipponhyoronsha, Tokyo, Japan [in Japanese].