

QUATERNIONIC REFORMULATION OF MASSIVE ELECTRODYNAMICS

A. S. Rawat

Department of Physics
H. N. B. Garhwal University Pauri Campus,
Pauri -246001 (U.K.), India

Abstract

Starting with the idea of two four-potentials and taking the generalized charge, potential, field, current as complex quantities with real and imaginary parts of them as electric and magnetic constituents, the quaternionic formulation of generalized complex-electromagnetic field equations and generalized Proca-Maxwell's (GPM) equations have been established for massive dyons.

1. Introduction

The magnetic monopole is a hypothetical elementary particle that is an isolated magnet with only one magnetic pole with a net magnetic charge. P. A. M. Dirac [1] showed that if any magnetic monopoles exist in the universe, then all electric charge in the universe must be quantized (called Dirac quantization condition). The electric charge is, quantized, which is consistent with the existence of monopoles. The subject of magnetic monopoles [1, 2] has intrigued and fascinated the physics community dating back to the early twentieth century. The eight decades of twentieth century witnessed a rapid development of the group theory and gauge field theory to establish the theoretical existence of monopoles and to explain their group properties and symmetries. Keeping in mind t'Hooft's solutions [3] and the fact that despite the potential importance of monopoles, the formalism necessary to describe them has been clumsy and not manifestly

covariant, Rajput et.al. [4, 5] developed the self-consistent quantum field theory of generalized electromagnetic fields associated with dyons (particles carrying electric and magnetic charges). The analogy between linear gravitational and electromagnetic fields leads the asymmetry in Einstein's linear equation of gravity and suggests the existence of gravitational analogue of magnetic monopole [6]. On the other hand, there has been a revival in the formulation of natural laws so that there exists [7] fourdivision algebras consisting the algebra of real numbers, complex numbers, quaternions and octonions. All four algebra's are alternative with totally anti symmetric associators.

Quaternions [8, 9] were very first example of hyper complex numbers have been widely used [10, 11, 12] to the various applications of mathematics and physics. Since quaternion is non commutative norm division algebra, it has four dimensional structure which may be useful to express the 4-dimensional theory of dyons. In order to understand the theoretical existence of monopoles as well as dyons in grand unified theories (GUTs) and keeping in view their recent potential importance, the fact that the formalism necessary to describe them has been clumsy, in the present paper we discuss the generalized electromagnetic EM-field equations of massive dyons along with the quaternion formulation in consistent manner. The quaternionic algebra shows the four dimensional representation of space-time in Euclidean space. We describe the generalized charge, potential, field, current source

of dyons in quaternionic form along with real and imaginary parts of electric and magnetic constituents. After that the quaternionic representation of generalized complex-electromagnetic (C-EM) field equations and generalized Proca-Maxwell's (GPM) equations for massive dyons have been reformulated.

2. Quaternion Preliminaries:

The algebra \mathbb{Q} of quaternion [8, 9] is a four-dimensional algebra over the field of real numbers \mathbb{R} . The quaternion is very suitable to express the typical four-qualities in physics, such as four-position, four-momentum, four-force, four-potential and four-current etc. Similar to a complex number $\mathbb{C} = (q_0 + iq_1)$ where $i^2 = -1$, a real quaternion \mathbb{Q} which is a four component number can be defined as

$$\mathbb{Q} = \sum_{j=0}^3 q_j e_j = \underbrace{q_0 e_0 + q_1 e_1 + q_2 e_2 + q_3 e_3}, \quad (2.1)$$

where the scalar part of the quaternion is represented by q_0 , the vector part is represented by q_1, q_2 and q_3 , while $e_0 = 1, e_j (j = 1, 2, 3)$ are quaternion units. The quaternion units (e_0, e_1, e_2, e_3) is known as the quaternion basis and its elements satisfy the following conditions,

$$\begin{aligned} e_0^2 &= e_0 = 1, e_j^2 = -e_0, \\ e_0 e_j &= e_j e_0 = e_j \quad (j = 1, 2, 3), \\ e_i e_j &= -\delta_{ij} + \varepsilon_{ijk} e_k \quad (\forall i, j, k = 1, 2, 3). \end{aligned} \quad (2.2)$$

Here δ_{ij} is the delta symbol and ε_{ijk} is the Levi Civita three index symbol having value $(\varepsilon_{ijk} = +1)$ for cyclic permutation, $(\varepsilon_{ijk} = -1)$ for anti cyclic permutation and $(\varepsilon_{ijk} = 0)$ for any two repeated indices. Addition and multiplication are defined by the usual distribution law $(e_j e_k) e_l = e_j (e_k e_l)$ along with the

multiplication rules given by equation (2.2). \mathbb{Q} is an associative but non commutative algebra. If q_0, q_1, q_2, q_3 are taken as complex quantities, the quaternion is said to be a bi-quaternion or complex quaternion. The quaternion conjugate $\bar{\mathbb{Q}} \{ e_0 \rightarrow e_0, e_j \rightarrow e_j (j = 1, 2, 3) \}$ is defined as

$$\bar{\mathbb{Q}} = \sum_{j=0}^3 q_j \bar{e}_j = \underbrace{q_0 e_0 - q_1 e_1 - q_2 e_2 - q_3 e_3}. \quad (2.3)$$

In practice \mathbb{Q} is often represented as a 2×2 matrix $\mathbb{Q} = q_0 - i\vec{\sigma} \cdot \vec{q}$ where $e_0 = I_{2 \times 2}$, $e_j = -i\sigma_j (j = 1, 2, 3)$ and σ_j are the usual Pauli spin matrices. The quaternion \mathbb{Q} and quaternion conjugate $\bar{\mathbb{Q}}$ may be written as a linear combination of scalar q_0 and spatial vector \vec{q} , i.e.

$$\begin{cases} \mathbb{Q} = (q_0, \vec{q}) = q_0 + (q_1 e_1 + q_2 e_2 + q_3 e_3), \\ \bar{\mathbb{Q}} = (q_0, -\vec{q}) = q_0 - (q_1 e_1 + q_2 e_2 + q_3 e_3). \end{cases} \quad (2.4)$$

The real part of the quaternion q_0 is defined as

$$\Re \mathbb{Q} = \frac{1}{2} (\mathbb{Q} + \bar{\mathbb{Q}}) = q_0, \quad (2.5)$$

where \Re denotes the real part and if $\Re \mathbb{Q} = 0$ then we have $\bar{\mathbb{Q}} = -\mathbb{Q}$ and imaginary \mathbb{Q} is known as pure quaternion written as

$$\mathbb{Q} = (q_1 e_1 + q_2 e_2 + q_3 e_3). \quad (2.6)$$

The quaternion product is associative and distributive:

$$\mathbb{P}(\mathbb{Q}\mathbb{R}) = (\mathbb{P}\mathbb{Q})\mathbb{R}, \mathbb{P}(\mathbb{Q} + \mathbb{R}) = \mathbb{P}\mathbb{Q} + \mathbb{P}\mathbb{R}. \quad (2.7)$$

The norm $N(\mathbb{Q})$ of a quaternion \mathbb{Q} may be expressed as

$$N(\mathbb{Q}) = \mathbb{Q}\bar{\mathbb{Q}} = \bar{\mathbb{Q}}\mathbb{Q} = \sum_{\alpha=0}^3 Q_\alpha^2 \cdot e_0. \quad (2.8)$$

Accordingly, the inverse of a quaternion \mathbb{Q} (non-zero norm) is defined as

$$\mathbb{Q}^{-1} = \frac{\bar{\mathbb{Q}}}{N(\mathbb{Q})} \Rightarrow \mathbb{Q}\mathbb{Q}^{-1} = \mathbb{Q}^{-1}\mathbb{Q} = 1. \quad (2.9)$$

3. Dyons (4-Dimensional Theory):

A dyon is defined as a particle which simultaneously carries electric and magnetic charge in 4-dimensional theory. The grand unified theories predicted the existence of both magnetic monopoles and dyons. Starting with the idea of Cabbing and Ferrari [13], the generalized charge, generalized four-potential, and generalized current of dyons can be written in terms of complex quantity with their real and imaginary parts of electric and magnetic constituents

$$\begin{cases} \mathcal{D} = e + ig; \\ \mathbf{V}^\nu = A^\nu + iB^\nu, \\ \mathbf{J}^\nu = \mathbf{J}^\nu + i\mathbf{K}^\nu, (\nu = 0, 1, 2, 3, \& i = \sqrt{-1}) \end{cases} \quad (3.1)$$

where (e, g; $A^\nu, B^\nu, \mathbf{J}^\nu$ and \mathbf{K}^ν) are respectively the electric charge, magnetic charge, electric four-potential, magnetic four-potential, electric four-current and magnetic four-current. We may write the symmetric Maxwell's equations known as Generalized Dirac Maxwell's (GDM) equations for dyons given by

$$\begin{aligned} \nabla \cdot \vec{E} &= \rho_e, \\ \nabla \cdot \vec{H} &= \rho_m, \\ \nabla \times \vec{E} &= -\vec{k} - \frac{\partial \vec{H}}{\partial t}, \\ \nabla \times \vec{H} &= -\vec{J} - \frac{\partial \vec{E}}{\partial t}. \end{aligned} \quad (3.2)$$

Here \vec{E} is the electric field, \vec{H} is the magnetic field, ρ_e is the charge source density due to electric charge, ρ_m is the charge source density due to magnetic charge (monopole), \vec{J} is the current source density due to electric charge and \vec{k} is the current source density due to magnetic charge. We have used the following notations:

$$\begin{cases} \{A^\nu\} = \{\vec{A}, \phi\}; \\ \{B^\nu\} = \{\vec{B}, \varphi\}; \\ \{J^\nu\} = \{\vec{J}, \rho_e\}; \\ \{K^\nu\} = \{\vec{K}, \rho_m\}. \end{cases} \quad (3.3)$$

We have also used the unit value of coefficients along with natural units ($c = \hbar = 1$) through out the text. The electric and magnetic fields of dyons satisfying the GDM equations (3.2) are now expressed in terms of components of two four potentials in a symmetrical manner i.e.

$$\begin{aligned} \vec{E} &= -\left(\nabla\phi + \frac{\partial \vec{A}}{\partial t} + \nabla \times \vec{B}\right), \\ \vec{H} &= -\left(\nabla\varphi + \frac{\partial \vec{B}}{\partial t} - \nabla \times \vec{A}\right). \end{aligned} \quad (3.4)$$

The complex vector field $\vec{\Psi}$ associated with generalized electromagnetic fields of dyons is defined as

$$\vec{\Psi} = (\vec{E} + i\vec{H}), \quad (3.5)$$

which reduces the four GDM equations (3.2) to following two differential equations as

$$\begin{aligned} \nabla \cdot \vec{\Psi} &= \zeta; \\ \nabla \times \vec{\Psi} &= -i\left(\vec{J} + \frac{\partial \vec{\Psi}}{\partial t}\right); \end{aligned} \quad (3.6)$$

where $\zeta = (\rho_e + i\rho_m)$ is the generalized charge density and $\vec{J} = (\vec{J} + i\vec{k})$ is the generalized current source density of dyons. These are regarded as the temporal and spatial components of generalized

four current $\{J^\nu\} = \{\vec{J}, \zeta\}$ of dyons. As such, we may write the generalized electromagnetic field vector $\vec{\Psi}$ in the following manner in terms of components of generalized four potential $\{V^\nu\} = \{\vec{V}, \Theta\}$ with $\Theta = (\varphi + i\phi)$ and $\vec{V} = (\vec{A} + i\vec{B})$ as

$$\vec{\Psi} = -\frac{\partial \vec{V}}{\partial t} - \nabla\Theta - i(\nabla \times \vec{V}). \quad (3.7)$$

Thus, we can express the GDM equations (3.2) of dyons in the following covariant notation [14],

$$\begin{aligned} F_{\mu\nu,\nu} &= \partial^\nu F_{\mu\nu} = J_\mu; \\ \mathcal{F}_{\mu\nu,\nu} &= \partial^\nu \mathcal{F}_{\mu\nu} = K_\mu; \end{aligned} \quad (3.8)$$

where $F_{0i} = E^i$ ($\nabla i = 1, 2, 3$),
 $F_{ij} = \varepsilon_{ijk} H^k$ ($\nabla i, j, k = 1, 2, 3$),
 $\mathcal{F}_{\mu\nu} = \frac{1}{2} \varepsilon_{\mu\nu\sigma\lambda} F^{\sigma\lambda}$ while $\varepsilon_{\mu\nu\sigma\lambda}$ is completely antisymmetric Ricci tensor of rank four. Therefore, we may write the Lagrangian which follows the minimum action principle for generalized electromagnetic fields of massive dyons as

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{4} \mathcal{F}_{\mu\nu} \mathcal{F}^{\mu\nu} + \frac{1}{2} \mu^2 A_\mu A^\mu \\ & + \frac{1}{2} \mu^2 B_\mu B^\mu - A_\mu J^\mu - B_\mu K^\mu, \end{aligned} \quad (3.9)$$

where μ^2 is the mass term of dyons. The Euler-Lagrange equations can be expressed as

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial A_\mu} - \partial_\nu \left(\frac{\partial \mathcal{L}}{\partial (\partial_\nu A_\mu)} \right) &= 0, \\ \frac{\partial \mathcal{L}}{\partial B_\mu} - \partial_\nu \left(\frac{\partial \mathcal{L}}{\partial (\partial_\nu B_\mu)} \right) &= 0. \end{aligned} \quad (3.10)$$

Then, the generalized Proca equations are defined:

$$\begin{aligned} \partial_\mu F^{\mu\nu} - \mu^2 A^\nu &= J^\nu; \\ \partial_\mu \mathcal{F}^{\mu\nu} - \mu^2 B^\nu &= K^\nu. \end{aligned} \quad (3.11)$$

The Proca equation [15] describing electrodynamics with finite range (or equivalently with a non-zero mass). In the case of massive dyons, the generalized fields equation may be written as [16],

$$\partial_\mu \mathbb{F}^{\mu\nu} - \mu^2 \mathbb{V}^\nu = \mathbb{J}^\nu. \quad (3.12)$$

On the other hand, the 't Hooft-Polyakov monopole solutions [3, 17, 18] describe an object

of finite size which from far away can not be distinguished from a Dirac monopole of charge $g \mapsto -\frac{1}{e}$. In view of the energy density of generalized electromagnetic-vector field, we may write the mass formula for dyon as

$$\mu(e, g) = \mu |e + ig| = \mu \sqrt{e^2 + g^2}, \quad (3.13)$$

which plays an important role in supersymmetric gauge theories. Furthermore, according to Bogomolny bound [22], the mass of a dyons (or massive dyon) is expressed as

$$\mu(e, g) = v |\mathcal{D}| = v |e + ig| = v \sqrt{e^2 + g^2},$$

which is known as BPS [19, 20] mass formula where v specifies the magnitude of the vacuum expectation value of scalar Higgs field. The BPS mass formula is universal and is also invariant under electromagnetic duality transformations. Thus, for massive magnetic monopole ($e \rightarrow 0$) or 't Hooft-Polyakov monopole, the Bogomolny bound [19] is given by

$$\mu(0, g) \geq v |g|, \quad (3.14)$$

which is possible in Prasad-Sommerfield limit [20].

4. Quaternion Reformulation of Massive Dyons:

Let us define the space time four differential operator as

$$\begin{aligned} \mathcal{D} = (\vec{\nabla} e_j, -i\partial_t e_0) &= \partial_x e_1, \partial_y e_2, \partial_z e_3, -i\partial_t e_0 \\ &= \partial_x e_1 + \partial_y e_2 + \partial_z e_3 + (-i\partial_t) e_0, \end{aligned} \quad (4.1)$$

which shows the quaternionic form of four differential operator

$$\left\{ \partial_\nu \rightarrow \frac{\partial}{\partial \nu} \right\} \Rightarrow (\vec{\nabla}, -i\partial_t),$$

and $\nu = 0, 1, 2, 3$. The quaternionic conjugate differential operator can be written as:

$$\begin{aligned} \bar{\mathcal{D}} &= (-\vec{\nabla}, -i\partial_t) \\ &= -\partial_x e_1 + \partial_y e_2 + \partial_z e_3 + (-i\partial_t) e_0. \end{aligned} \quad (4.2)$$

Then, the product of $\mathcal{D}\bar{\mathcal{D}}$ will be

$$\begin{aligned} \mathcal{D}\bar{\mathcal{D}} &= (\partial_x^2 + \partial_y^2 + \partial_z^2 - \partial_t^2) \\ &= (\nabla^2 - \partial_t^2) = \bar{\mathcal{D}}\mathcal{D}. \end{aligned} \quad (4.3)$$

Here $(\nabla^2 - \partial_t^2) = \square$ is defined the D' Alembert operator. In order to consider the generalized electromagnetic fields of massive dyons, we can define quaternionic form of generalized four potential of dyons as

$$\mathbf{V} = (\vec{V}, \Theta) \equiv \underbrace{V_x e_1 + V_y e_2 + V_z e_3 + \Theta e_0}_{\mathbf{V}_\mu}, \quad (4.4)$$

where $(V_x, V_y, V_z, \Theta) = (\vec{V}, \Theta) = \{\mathbf{V}_\mu\}$ are described as the components of generalized four potential associated with generalized charge $(\mathcal{D} = e + ig)$ of dyons. We have assumed the generalized dyonic charge (\mathcal{D}) which is complex in nature [5], as the same way we can take the

quaternionic generalized potential (\mathbf{V}_μ) of dyons in complex nature. Thus, we may now identify the components of generalized complex-potential $\mathbf{V}_\mu \rightarrow (A_\mu + iB_\mu)$ of massive dyons as

$$\begin{aligned} V_x &\mapsto (A_x + iB_x), V_y \mapsto (A_y + iB_y), \\ V_z &\mapsto (A_z + iB_z), \Theta \mapsto (\varphi + i\phi). \end{aligned} \quad (4.5)$$

Therefore, the generalized complex-potential $(\mathbf{V} \mapsto \mathbf{V}_\mathbb{C})$ of massive dyons can be expressed in terms of quaternionic elements,

$$\begin{aligned} \mathbf{V}_\mathbb{C} = (\vec{V}, \Theta) &\equiv (A_x + iB_x) e_1 + (A_y + iB_y) e_2 \\ &+ (A_z + iB_z) e_3 + (\varphi + i\phi) e_0. \end{aligned} \quad (4.6)$$

As such, we can write the generalized quaternionic electromagnetic fields Ψ of massive dyons as

$$\mathcal{D}\mathbf{V}_\mathbb{C} = \Psi \mapsto (\vec{\psi}, \chi_t), \quad (4.7)$$

where Ψ is defined by

$$\Psi = (\vec{\psi}, \chi_t) \equiv \underbrace{\Psi_x e_1 + \Psi_y e_2 + \Psi_z e_3 + \chi_t e_0}_{\Psi_\mathbb{C}} \quad (4.8)$$

In quaternionic representation of Y, we have used the EM-field components

$$\vec{\psi} \{ \vec{H}, \vec{E} \} = (\psi_x, \psi_y, \psi_z) \text{ and } \chi_t \{ H_0, E_0 \} = 0$$

because of Lorenz gauge conditions applied separately on electric and magnetic four potential. Using (4.7), the generalized quaternionic complex-electromagnetic fields $(\Psi \mapsto \Psi_\mathbb{C})$ of dyons can be written as

$$\Psi_\mathbb{C} = (H_x + iE_x) e_1 + (H_y + iE_y) e_2 + (H_z + iE_z) e_3, \quad (4.9)$$

where

$\psi_x \rightarrow (H_x + iE_x), \psi_y \rightarrow (H_y + iE_y), \psi_z \rightarrow (H_z + iE_z)$ are respectively the components of complex-electromagnetic field for massive dyons. Thus the generalized vector field can be expressed in terms of following quaternionic form,

$$\begin{aligned} \Psi_\mathbb{C} &\mapsto (\vec{H} + i\vec{E}) e_j \\ &= \left(-\vec{\nabla} \phi - \frac{\partial \vec{B}}{\partial t} + \vec{\nabla} \times \vec{A} \right) e_j + i \left(-\vec{\nabla} \phi - \frac{\partial \vec{A}}{\partial t} - \vec{\nabla} \times \vec{B} \right) e_j, \end{aligned}$$

where the electric and magnetic field vectors for dyons are already defined in equation (3.4).

4.1. Quaternion Reformulation Generalized Proca-Maxwell's (GPM) Equations for Massive Dyons:

In order to obtain the generalized massive field equations of dyons in four-dimensional space-

time, we may now applying differential operator \mathcal{D} to the generalized quaternion complex EM-fields $\Psi_{\mathbf{C}}$ given by the following way

$$\begin{aligned} \mathcal{D}\Psi_{\mathbf{C}} = & \left\{ \left[(\vec{\nabla} \times \vec{H})_x - \frac{\partial E_x}{\partial t} \right] + i \left[(\vec{\nabla} \times \vec{E})_x - \frac{\partial H_x}{\partial t} \right] \right\} e_1 \\ & + \left\{ \left[(\vec{\nabla} \times \vec{H})_y - \frac{\partial E_y}{\partial t} \right] + i \left[(\vec{\nabla} \times \vec{E})_y - \frac{\partial H_y}{\partial t} \right] \right\} e_2 \\ & + \left\{ \left[(\vec{\nabla} \times \vec{H})_z - \frac{\partial E_z}{\partial t} \right] + i \left[(\vec{\nabla} \times \vec{E})_z - \frac{\partial H_z}{\partial t} \right] \right\} e_3 \\ & + \left\{ [(\vec{\nabla} \cdot \vec{H}) + i(\vec{\nabla} \cdot \vec{E})] \right\} e_0. \end{aligned} \quad (4.10)$$

The definition for the general quaternionic field of massive dyons will be,

$$\mathcal{D}\Psi_{\mathbf{C}} = \left(\mathbb{J}_{\mathbf{C}} + \mu^2 \mathbb{V}_{\mathbf{C}} \right), \quad (4.11)$$

where μ^2 is the mass term for dyons given by (2.13) and $\mathbb{J}_{\mathbf{C}}$ is the quaternionic form of generalized complex-current source of dyons. Thus using generalized complex potential of dyons, we can write

$$\begin{aligned} \mu^2 \mathbb{V}_{\mathbf{C}} = & \mu^2 (A_x + iB_x) e_1 + \mu^2 (A_y + iB_y) e_2 \\ & + \mu^2 (A_z + iB_z) e_3 + \mu^2 (\varphi + i\phi) e_0, \end{aligned} \quad (4.12)$$

and the quaternionic form of generalized complex-current

$$\mathbb{J}_{\mathbf{C}} = (\vec{J}, \zeta) = \underbrace{J_x e_1 + J_y e_2 + J_z e_3 + \zeta e_0}_{(4.13)},$$

which may be reduced in term of dyonic form, i.e.,

$$\begin{aligned} \mathbb{J}_{\mathbf{C}} &= \underbrace{\sum_{n=1}^3 (\vec{j} + i\vec{k}) e_n}_{(Current\ density\ for\ Dyons)} + \underbrace{(\rho_m + i\rho_e) e_0}_{(Charge\ density\ for\ Dyons)} \\ &= (j_x + ik_x) e_1 + (j_y + ik_y) e_2 \\ &+ (j_z + ik_z) e_3 + (\rho_m + i\rho_e) e_0. \end{aligned} \quad (4.14)$$

here $(\vec{j}, \rho_e) = \{J_{\mu}\}, (\vec{k}, \rho_m) = \{K_{\mu}\}$ and $(\vec{J}, \zeta) = \{\mathbb{J}_{\mu}\}$ are respectively the four currents associated with electric charge, magnetic monopole and generalized fields of massive dyons. Thus the right hand part of the equation (4.11) can be expressed as

$$\begin{aligned} & \left(\mathbb{J}_{\mathbf{C}} + \mu^2 \mathbb{V}_{\mathbf{C}} \right) \\ &= \sum_{n=1}^3 \left[(j_n + ik_n) + \mu^2 (A_n + iB_n) \right] e_n \\ &+ \left[(\rho_m + i\rho_e) + \mu^2 (\varphi + i\phi) \right] e_0. \end{aligned} \quad (4.15)$$

From equation (4.11), we get the following differential equations

$$\begin{aligned} & \left[(\vec{\nabla} \times \vec{H})_x - \frac{\partial E_x}{\partial t} - \mu^2 A_x \right] = j_x; (\text{vector term of } e_1) \\ & \left[(\vec{\nabla} \times \vec{H})_y - \frac{\partial E_y}{\partial t} - \mu^2 A_y \right] = j_y; (\text{vector term of } e_2) \\ & \left[(\vec{\nabla} \times \vec{H})_z - \frac{\partial E_z}{\partial t} - \mu^2 A_z \right] = j_z; (\text{vector term of } e_3) \\ & \left[(\vec{\nabla} \times \vec{E})_x + \frac{\partial H_x}{\partial t} + \mu^2 B_x \right] = -k_x; (\text{imaginary vector term of } e_1) \\ & \left[(\vec{\nabla} \times \vec{E})_y + \frac{\partial H_y}{\partial t} + \mu^2 B_y \right] = -k_y; (\text{imaginary vector term of } e_2) \\ & \left[(\vec{\nabla} \times \vec{E})_z + \frac{\partial H_z}{\partial t} + \mu^2 B_z \right] = -k_z; (\text{imaginary vector term of } e_3) \\ & \left[(\vec{\nabla} \cdot \vec{H}) + \mu^2 \varphi \right] = \rho_m; (\text{scalar term of } e_0) \\ & \left[(\vec{\nabla} \cdot \vec{E}) - \mu^2 \phi \right] = \rho_e; (\text{imaginary scalar term of } e_0) \end{aligned} \quad (4.16)$$

Thus the quaternionic formulation can be used to study each components of generalized EM-field of massive dyons. As such, equation (4.16) leads to following compact form:

$$\begin{aligned}
 (\vec{\nabla} \cdot \vec{H}) + \mu^2 \phi &= \rho_m; \\
 (\vec{\nabla} \cdot \vec{E}) - \mu^2 \phi &= \rho_e; \\
 (\vec{\nabla} \times \vec{H}) - \mu^2 \vec{A} &= \frac{\partial \vec{E}}{\partial t} + \vec{j}; \\
 (\vec{\nabla} \times \vec{E}) + \mu^2 \vec{B} &= -\frac{\partial \vec{H}}{\partial t} - \vec{k};
 \end{aligned}
 \tag{4.17}$$

which are the generalized Proca-Maxwell's (GPM) equations of generalized fields of massive dyons. If we put $\mu \rightarrow 0$ i.e. massless particles, then (4.17) exhibits the simplest generalized Dirac-Maxwell's equation for dyons. The present quaternion reformulation of generalized fields of massive dyons represents well the invariance of field equations under Lorentz and duality transformations. It also reproduces the dynamics of electric (magnetic) charge yielding to the usual form of Maxwell's equations in the absence of magnetic (electric charge) in compact, simpler and consistent way.

5. Conclusion

In this paper, the electric and magnetic four complex-potentials of dyons have been considered in usual four-dimensional quaternionic space. In four-dimensional quaternionic theory, a massive dyon is a hypothetical particle with both electric and magnetic charges. A massive dyon with a zero electric charge is usually referred to as a massive magnetic monopole. Thus, the quaternionic formulation of generalized complex-electromagnetic fields equations and generalized Proca-Maxwell's (GPM) equations of massive dyons have been described. Equation (4.17) is the compact form of GPM equations of massive dyons. Hence, we concluded that the four dimensional quaternion algebra is very simple and consistent to represent the electromagnetic theory of massive dyons

6. References.

- [1] P. A. M. Dirac, "Quantized singularities in the electromagnetic field", Proc. Roy. Soc. London, A133 (1931), 60.
- [2] J. Schwinger, "Dyons Versus Quarks", Science, 166 (1969), 690.
- [3] G. pt Hooft, "Magnetic monopoles in unified gauge theories", Nucl. Phys., B79 (1974), 276.
- [4] P. S. Bisht, O. P. S. Negi and B. S. Rajput, "Null-tetrad formulation of dyons", IL Nuovo Cimento 104A (1991), 337.
- [5] B. S. Rajput, S. R. Kumar and O. P. S. Negi, "Quaternionic formulation for dyons", Lett. Nuovo Cimento, 34 (1982), 180.
- [6] J. S. Dowker and J. A. Roche, "The Gravitational Analogues of Magnetic Monopoles", Proc. R. Soc., 92 (1967), 1.
- [7] L. E. Dickson, "On Quaternions and Their Generalization and the History of the Eight Square Theorem", Ann. Math., 20 (1919), 155.
- [8] W. R. Hamilton, "Elements of quaternions", Chelsea Publications Co., NY, (1969).
- [9] P. G. Tait, "An elementary Treatise on Quaternions", Oxford Univ. Press, NY, (1875).
- [10] D. Finklestein, J. M. Jauch, S. Schiminovich and D. Speiser, "Principle of general quaternion covariance", J. Math. Phys., 4 (1963), 788.

- [11] S. L. Adler, "Quaternion Quantum Mechanics and Quantum Fields", Oxford Univ. Press, NY, (1995).
- [12] P. S. Bisht, O. P. S. Negi and B. S. Rajput, "Quaternion Gauge Theory of Dyon Fields" Prog. Theor. Phys., 85 (1991), 157.
- [13] N. Cabibbo and E. Ferrari, "Quantum electrodynamics with Dirac monopoles", Nuovo Cim., 23 (1962), 1147.
- [14] B. S. Rajput, S. Kumar and O. P. S. Negi, "Quaternionic formulation for generalized field equations in the presence of dyons", Lett. Nuovo Cimento, 34 (1982), 180.
- [15] A. Proca, "Sur l'equation de Dirac" Compt. Rend. 190 (1930), 1377.
- [16] A.Yu. Ignatiev and G. C. Joshi, "Massive electrodynamics and the magnetic monopoles", Phys. Rev., D 53 (1996), 984.
- [17] G. 't Hooft, "A property of electric and magnetic flux in non-Abelian gauge theories", Nucl. Phys., B153 (1979), 141.
- [18] A. M. Polyakov, "Particle spectrum in the quantum field theory", JETP. Lett., 20 (1974), 194.
- [19] E. B. Bogomolny, "Stability of Classical Solutions", Sov. J. Nucl. Phys., 24 (1976), 449.
- [20] M. K. Prasad and C. M. Sommerfield, "Exact Classical Solution for the 't Hooft Monopole and the Julia-Zee Dyon", Phys. Rev. Lett., 35 (1975), 760.