

Magnetohydrodynamic Boundary Layer Flow of Williamson Nanofluid over a Moving Surface in the presence of Gyrotactic Microorganism

Rekha K.¹, Asha C. S.² and Achala L. Nargund³,

^{1,2} and ³ P. G. Dept of Mathematics and Research Centre in Applied Mathematics,
M. E. S. College of Arts, Commerce and Science, Malleswaram, Bangalore-560003.

Abstract

The two-dimensional steady incompressible fluid flow of MHD Williamson nanofluid with gyrotactic microorganisms over a moving surface is considered. The four nonlinear governing partial differential equations are reduced to ordinary differential equations, by using a similarity transformation. Homotopy Analysis Method (HAM) is applied to solve the reduced equations and the effect of importance of fluid parameters are explained through graphs. Numerical solution is also obtained and shown that HAM solution matches exactly. The radius of convergence of the solution is determined through Domb-Sykes plot.

Keywords: Williamson nanofluid, Brownian motion, Gyrotactic Microorganisms, MHD, Domb-Sykes plot.

1 INTRODUCTION

Nanofluids are made up of nanoparticles (diameter of less than 100 nanometers) suspended in a base fluid such as water, oil, or ethylene glycol. Nanofluids offer a large surface area for heat transmission between particles and fluids. Nanofluids are characterized by their thermal conductivity. Nanofluids offer significant benefits in a variety of biomedical applications, including cancer therapy, safer surgery by cooling, high-powered lasers, X-ray generators, and magnetic cell separation.

Bioconvection is an occurrence that is characterized by the movement of upward swimming microorganisms such as *Bacillus subtilis*, *Chlamydomonas nivalis*, *Oxyatic*

bacteria etc. The nanofluids microorganisms such as oxytactic gyrotactic microorganisms. Here we considered the gyrotactic microorganisms which are very tiny, single celled, self propelled and locally occurring. Due the small scale of microorganisms inertial effect can be neglected and the stability of the nanoparticle suspension may increase. Such type of gyrotactic microorganisms exist in fluids such as seas, lakes, streams and water storages.

Nanoparticles containing microorganisms can improve the thermal efficiency of several systems, including bacteria-powered micro-mixers, microfluidics devices like micro-volumes and enzyme biosensors, microbial fuel cells and bio-microsystems like chip-shaped microdevices. Mobile microbes can enhance the stability of nanoparticle suspensions, microscale mixing, enhanced mass transfer and microvolume use of the suspensions [1, 2, 3, 4]. The development of patterns in layered suspensions of negatively geotactic microbes are discussed by Childress et al. [5].

Talha et al. [6] discussed the two-dimensional steady magnetohydrodynamic boundary layer flow of Williamson nanofluid containing gyrotactic microorganism in the presence of exponential internal heat generation over a stretching sheet and compared the results with and without exponential internal heat generation using Spectral Relaxation Technique (SRT) for numerical solution. Recently Alharbi et al. [7] investigated the two-dimensional stable electrically conducting hybrid nanofluid laminar mixed convection incompressible flow with viscous and gyrotactic microorganism near the stagnation point. A theoretical analysis of migrating gyrotactic microorganism swimming through an isotropically narrowing artery in a non-Newtonian blood base nanofluid is studied by Bhatti et al. [8].

Al-Khaled et al. [9] analyzed the unsteady incompressible fluid flow of tangent hyperbolic nanofluid over an accelerated moving surface and used HAM to find the analytical solution. Lv et al. [10] studied the steady flow of hybrid nanofluid over a spinning disk and used the Parametric Continuation Method (PCM) to solve the transformed equations. The two-dimensional MHD heat transfer and boundary layer flow of Williamson fluid on the exponentially vertical shrinking sheet with variable thickness and thermal conductivity are analyzed by Lund et al. [11]. Atif et al. [12] studied the MHD stratified micropolar bioconvective flow of micropolar nanofluid with gyrotactic microorganisms and converted system of PDE's are solved by Shooting technique. Yusuf et al. [13] studied the bioconvective flow of Williamson nanofluid under the impact of heat radiation and gyrotactic microorganism in a porous of an aligned semi-infinite plate.

Shahid et al. [14] investigated the effect of the magnetic field, thermal radiation and chemical reaction of nanofluid flow using Successive Taylor's Series Linearization Method (STSLM). The flow of nanofluid over an isothermal vertical porous Wedge containing gyrotactic microorganisms using the Keller Box method is studied by Mahdy et al. [15]. Non-linear radiation and variable viscosity in two different cases such as oblique and free stream flow are considered by Jayachandra Babu et al. [16] and analyzed the bio conductive stagnation point flow of Williamson nanofluid across a stretching sheet. Khan et al. [17] analyzed the combined impacts of radiative heat

and variable viscosity on the oblique hydromagnetic flow with mass and heat transfer of nanofluid over a stretching boundary layer. The bioconvection of a nanofluid with the effects of magnetic field and convective boundary state is examined by Chakrabarty et al. [18]. Dual solutions for the flow of Casson fluid over a vertical rotating cone in a porous medium is investigated by Raju et al. [19].

Bees et al. [20] proposed the nonlinear structure of deep, stochastic, gyrotactic bioconvection. A linear analysis is reviewed and a weakly non-linear analysis is used to show the supercritical nature of the bifurcation. Akbar [21] proposed a peristaltic flow of a nanofluid in an asymmetric channel including gyrotactic microorganism and provided numerical results using long wavelength and low Reynold's number approximation. A dilute suspension of gyrotactic and oxytactic microorganisms in water is proposed by Kuznetsov [22, 23]. He concluded that gyrotactic microorganisms always have a destabilizing effect and the Galerkin method is applied to get the results. Ahmed et al. [24, 25] explained the flow of nanofluids across a porous media and obtained numerical solution by using the Successive over Relaxation (SOR) parameter method.

Ramzan et al. [26] concluded that the effect of increase in bioconvection Lewis number and Peclet number of microorganisms decreases the motile density. The effects of solar radiation on hydromagnetic bioconvection of water-based nanofluid flow via a permeable surface in the residence of gyrotactic microorganisms are examined by Acharya et al. [27]. Achala et al. [37, 38, 39, 40] applied HAM to solve ordinary and partial differential equations, obtained region of convergence by Domb-Sykes plot and applied the Pade approximation to identify the singularities. Sathyanarayana et al. [41, 42] applied HAM to obtain exact analytical solution to the two dimensional laminar compressible boundary layer flow with pressure gradient.

In this paper we are analysing Williamson nanofluid model in boundary layer over a moving surface by considering presence of gyrotactic microorganisms. We are extending the work of Talha et al. [6] for moving surface. The distribution of the paper is first section contains Introduction, second section involves governing equations of the problem, third section is Homotopy Analysis solution of the problem, fourth section is result and discussion and fifth section consists of graphs and tables.

2 MATHEMATICAL FORMULATION

The two dimensional steady incompressible fluid flow of MHD Williamson nanofluid with gyrotactic microorganism over a moving surface [6] is considered. u and v are velocity components along x and y directions respectively. Uniform magnetic field B_0 is considered. The fundamental equations for this model are as follows.

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} = 0, \quad (1)$$

$$u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} = \nu \frac{\partial^2 u}{\partial y^2} + \sqrt{2} \nu \Gamma \frac{\partial u}{\partial y} \frac{\partial^2 u}{\partial y^2} - \frac{\sigma B_0^2 u}{\rho}, \quad (2)$$

$$u \frac{\partial T}{\partial x} + v \frac{\partial T}{\partial y} = \alpha \frac{\partial^2 T}{\partial y^2} + \frac{\rho_p c_p}{\rho c} \left(D_B \frac{\partial C}{\partial y} \frac{\partial T}{\partial y} + \frac{D_T}{T_\infty} \left(\frac{\partial T}{\partial y} \right)^2 \right), \quad (3)$$

$$u \frac{\partial C}{\partial x} + v \frac{\partial C}{\partial y} = D_B \frac{\partial^2 C}{\partial y^2} + \frac{D_T}{T_\infty} \frac{\partial^2 T}{\partial y^2}, \quad (4)$$

$$u \frac{\partial n}{\partial x} + v \frac{\partial n}{\partial y} = D_n \frac{\partial^2 n}{\partial y^2} - \frac{dW_c}{c_w - c_\infty} \frac{\partial}{\partial y} \left(n \frac{\partial C}{\partial y} \right), \quad (5)$$

where u and v are velocity components in x and y direction, ν is the kinematic viscosity, α is the thermal diffusivity of nanofluid, ρ is density of the fluid, B_0 is the magnetic field, σ is the electrical conductivity, τ is heat capacity ratio, T is the temperature of the fluid, C is the concentration of the fluid, D_T is thermophoresis coefficient, D_B is Brownian diffusion coefficient, n is the density of the motile microorganism, D_n is diffusivity of microorganism, W_c is maximum swimming speed, λ is power exponent parameter. The corresponding boundary conditions are given by

$$u = \lambda_1 U, v = 0, T = T_\omega = T_\infty + Ax^\lambda, C = C_\omega = C_\infty + Bx^\lambda, n = n_\omega \\ = n_\infty + Ex^\lambda \text{ at } y = 0;$$

$$u \rightarrow U, T \rightarrow T_\infty, C \rightarrow C_\infty, n \rightarrow n_\infty \text{ as } y \rightarrow \infty. \quad (6)$$

Introducing the following similarity transformations to reduce the governing equations into a system of ordinary differential equations.

$$\Psi = \sqrt{(2U\nu x)} f(\eta), \theta(\eta) = \frac{T - T_\infty}{T_\omega - T_\infty}, \phi(\eta) = \frac{C - C_\infty}{C_\omega - C_\infty}, \xi(\eta) = \frac{n - n_\infty}{n_\omega - n_\infty}, \eta = y \sqrt{\frac{U}{2\nu x}}. \quad (7)$$

Where Ψ is stream function. Substituting equation (7) in (2) - (6) we get

$$f'''(1 + \gamma f'') + f f'' - M f' = 0, \quad (8)$$

$$\theta'' + Pr(f + Nb\phi')\theta' - 2Pr\lambda f'\theta + PrNt(\theta')^2 = 0, \quad (9)$$

$$\phi'' + Lef\phi' - 2Le\lambda f'\phi + \frac{Nt}{Nb}\theta'' = 0, \quad (10)$$

$$\xi'' + (PrLbf - Pe\phi')\xi' - (2PrLb\lambda f' + Pe\phi'')\xi - Pe\sigma\phi'' = 0. \quad (11)$$

$$f = 0, f' = \lambda_1, \theta = 1, \phi = 1, \xi = 1 \text{ at } \eta = 0,$$

$$f' \rightarrow 1, \theta \rightarrow 0, \phi \rightarrow 0, \xi \rightarrow 0 \text{ as } \eta \rightarrow \infty. \quad (12)$$

Where

$$M = \left(\frac{2\sigma B_0^2 x}{\rho U} \right); \text{ Magnetic field parameter,}$$

$$\gamma = \Gamma \frac{U^{\frac{3}{2}}}{\sqrt{\nu x}}; \text{ Non Newtonian Williamson parameter,}$$

$$Pr = \frac{\nu}{\alpha}; \text{ Prandtl number,}$$

$$Pe = \left(\frac{dW_c}{D_n} \right); \text{ Peclet number,}$$

$$Le = \frac{\nu}{D_B}; \text{ Lewis number,}$$

$$Nb = \left(\frac{(\rho c_p) D_B (C_w - C_\infty)}{\nu (\rho c_f)} \right); \text{ Brownian motion parameter,}$$

$$Nt = \left(\frac{(\rho c_p) D_T (T_w - T_\infty)}{\nu T_\infty (\rho c_f)} \right); \text{ Thermophoresis parameter,}$$

$$Lb = \frac{\alpha}{D_n}; \text{ Bioconvection Lewis number,}$$

$$\sigma = \frac{n_\infty}{n_w - n_\infty}; \text{ Bioconvection parameter.}$$

3 HOMOTOPY ANALYSIS SOLUTION

Shijun Liao (1992) [29, 30, 31, 32, 33, 34, 35] explained Homotopy Analysis Method (HAM) to solve non-linear differential equations analytically. Using this method [36] we solve coupled nonlinear equations of this problem. The steps of the method are,

$$N[f(\eta)] = f'''(1 + \gamma f'') + ff'' - Mf', \quad (13)$$

$$N[\theta(\eta)] = \theta'' + Pr(f + Nb\phi')\theta' - 2Pr\lambda f'\theta + PrNt(\theta')^2, \quad (14)$$

$$N[\phi(\eta)] = \phi'' + Lef\phi' - 2Le\lambda f'\phi + \frac{Nt}{Nb}\theta'', \quad (15)$$

$$N[\xi(\eta)] = \xi'' + (PrLbf - Pe\phi')\xi' - (2PrLb\lambda f' + Pe\phi'')\xi - Pe\sigma\phi''. \quad (16)$$

Linear operators considered are as follows,

$$L_f(f) = \frac{\partial^3 f}{\partial \eta^3} + \frac{\partial^2 f}{\partial \eta^2}, \quad (17)$$

$$L_\theta(\theta) = \frac{\partial^2 \theta}{\partial \eta^2} + \frac{\partial \theta}{\partial \eta}, \quad (18)$$

$$L_\phi(\phi) = \frac{\partial^2 \phi}{\partial \eta^2} + \frac{\partial \phi}{\partial \eta}, \quad (19)$$

$$L_{\xi}(\xi) = \frac{\partial^2 \xi}{\partial \eta^2} + \frac{\partial \xi}{\partial \eta}, \quad (20)$$

which gives initial approximations as,

$$f_0 = (\lambda_1 - 1) + \eta + (1 - \lambda_1)e^{-\eta}, \quad (21)$$

$$\theta_0 = e^{-\eta}, \quad (22)$$

$$\phi_0 = e^{-\eta}, \quad (23)$$

$$\xi_0 = e^{-\eta}. \quad (24)$$

The nonlinear equations for approximate solutions are,

$$(1 - p)L_f[f(\eta, p) - f_0(\eta)] = hp \left[\frac{\partial^3 f}{\partial \eta^3} \left(1 + \gamma \frac{\partial^2 f}{\partial \eta^2} \right) + f \frac{\partial^2 f}{\partial \eta^2} - M \frac{\partial f}{\partial \eta} \right], \quad (25)$$

$$(1 - p)L_{\theta}[\theta(\eta, p) - \theta_0(\eta)] = hp \left[\frac{\partial^2 \theta}{\partial \eta^2} + Pr \left(f + Nb \frac{\partial \phi}{\partial \eta} \right) \frac{\partial \theta}{\partial \eta} - 2Pr\lambda \frac{\partial f}{\partial \eta} \theta + PrNt \left(\frac{\partial \theta}{\partial \eta} \right)^2 \right], \quad (26)$$

$$(1 - p)L_{\phi}[\phi(\eta, p) - \phi_0(\eta)] = hp \left[\frac{\partial^2 \phi}{\partial \eta^2} + Le f \frac{\partial \phi}{\partial \eta} - 2Le\lambda \frac{\partial f}{\partial \eta} \phi + \frac{Nt}{Nb} \frac{\partial^2 \theta}{\partial \eta^2} \right], \quad (27)$$

$$(1 - p)L_{\xi}[\xi(\eta, p) - \xi_0(\eta)] = hp \left[\frac{\partial^2 \xi}{\partial \eta^2} + \left(PrLbf - Pe \frac{\partial \phi}{\partial \eta} \right) \frac{\partial \xi}{\partial \eta} - \left(2PrLb\lambda \frac{\partial f}{\partial \eta} + Pe \frac{\partial^2 \phi}{\partial \eta^2} \right) \xi - Pe\sigma \frac{\partial^2 \phi}{\partial \eta^2} \right], \quad (28)$$

with following boundary conditions,

$$f(0, p) = 0, f_{\eta}(0, p) = \lambda_1, f_{\eta}(\infty, p) = 1, \quad (29)$$

$$\theta(0, p) = 1, \theta(\infty, p) = 0, \quad (30)$$

$$\phi(0, p) = 1, \phi(\infty, p) = 0, \quad (31)$$

$$\xi(0, p) = 1, \xi(\infty, p) = 0. \quad (32)$$

Varying the values of p from 0 to 1 we get the solution from first approximation to required solution. Defining

$$f_0(\eta) = f(\eta, 0) = \zeta_0(\eta), \quad (33)$$

$$\theta_0(\eta) = \theta(\eta, 0) = \mu_0(\eta), \quad (34)$$

$$\phi_0(\eta) = \phi(\eta, 0) = \psi_0(\eta), \quad (35)$$

$$\xi_0(\eta) = \xi(\eta, 0) = \tau_0(\eta). \quad (36)$$

Using Maclaurin's series expansion and applying Leibnitz theorem we get the series solution. The convergence of the series solution is derived by calculating the convergence parameter h .

$$L_f[\zeta_k - \chi_k \zeta_{k-1}] = hr_k(\eta), \quad (37)$$

$$L_\theta[\mu_k - \chi_k \mu_{k-1}] = hs_k(\eta), \quad (38)$$

$$L_\phi[\psi_k - \chi_k \psi_{k-1}] = ht_k(\eta), \quad (39)$$

$$L_\xi[\tau_k - \chi_k \tau_{k-1}] = hw_k(\eta). \quad (40)$$

$$\text{where } \chi_k = \begin{cases} 0, & \text{when } k \leq 1 \\ 1, & \text{when } k > 1 \end{cases} \quad \text{and} \quad (41)$$

$$r_k(\eta) = \zeta_{k-1}'''(\eta) + \gamma \sum_{m=0}^{k-1} \zeta_{k-1-m}''(\eta) \zeta_m'''(\eta) + \sum_{m=0}^{k-1} \zeta_{k-1-m}(\eta) \zeta_m''(\eta) - M \zeta_{k-1}'(\eta), \quad (42)$$

$$s_k(\eta) = \mu_{k-1}''(\eta) + Pr \left(\sum_{m=0}^{k-1} \zeta_{k-1-m}(\eta) + Nb \sum_{m=0}^{k-1} \psi_{k-1-m}'(\eta) \right) \mu_m'(\eta) - 2Pr\lambda \sum_{m=0}^{k-1} \zeta_{k-1-m}'(\eta) \mu_m(\eta) + PrNt \sum_{m=0}^{k-1} \mu_{k-1-m}'(\eta) \mu_m'(\eta), \quad (43)$$

$$t_k(\eta) = \psi_{k-1}''(\eta) + Le \sum_{m=0}^{k-1} \zeta_{k-1-m}(\eta) \psi_m'(\eta) - 2Le\lambda \sum_{m=0}^{k-1} \zeta_{k-1-m}'(\eta) \psi_m(\eta) + \frac{Nt}{Nb} \mu_{k-1}''(\eta), \quad (44)$$

$$w_k(\eta) = \tau_{k-1}''(\eta) + (PrLb \sum_{m=0}^{k-1} \zeta_{k-1-m}(\eta) - Pe \sum_{m=0}^{k-1} \psi_{k-1-m}'(\eta)) \tau_m'(\eta) - (2PrLb\lambda \sum_{m=0}^{k-1} \zeta_{k-1-m}'(\eta) + Pe \sum_{m=0}^{k-1} \psi_{k-1-m}''(\eta)) \tau_m(\eta) - Pe\sigma \psi_{k-1}''(\eta), \quad (45)$$

with boundary conditions,

$$\zeta_k(0) = 0, \zeta_k'(0) = 0, \zeta_k'(\infty) = 0, \quad (46)$$

$$\mu_k(0) = 0, \mu_k(\infty) = 0, \quad (47)$$

$$\psi_k(0) = 0, \psi_k(\infty) = 0, \quad (48)$$

$$\tau_k(0) = 0, \tau_k(\infty) = 0. \quad (49)$$

The required solution is

$$f = \zeta_0 + \zeta_1 + \zeta_2 + \zeta_3 + \dots, \quad (50)$$

$$\theta = \mu_0 + \mu_1 + \mu_2 + \mu_3 + \dots, \quad (51)$$

$$\phi = \psi_0 + \psi_1 + \psi_2 + \psi_3 + \dots, \quad (52)$$

$$\xi = \tau_0 + \tau_1 + \tau_2 + \tau_3 + \dots \quad (53)$$

Solving equations (37)-(40) by using MATHEMATICA we get

$$\begin{aligned} \zeta_1 = & \frac{1}{4}(-h - 4hM + h\gamma - 2h\lambda 1 + 4hM\lambda 1 - 2h\gamma\lambda 1 + 3h\lambda 1^2 + h\gamma\lambda 1^2) + \frac{1}{4}e^{-2\eta} \\ & \left(2e^\eta(3h - 2hM - h\gamma - 6h\lambda 1 + 2hM\lambda 1 + 2h\gamma\lambda 1 + 3h\lambda 1^2 - h\gamma\lambda 1^2) + \right. \\ & \left. h(-2e^{2\eta}M\eta^2 + (-1 + \gamma)(-1 + \lambda 1)^2 - 2e^\eta(-1 + \lambda 1)(-2 + \eta^2 + 2M(2 + \eta) + \right. \\ & \left. 4\lambda 1 + 2\eta\lambda 1)) \right), \dots \end{aligned} \quad (54)$$

$$\begin{aligned} \mu_1 = & \frac{1}{2}e^{-2\eta} \left(hPr(-1 + Nb + Nt + 2\lambda + \lambda 1 - 2\lambda\lambda 1) + e^\eta \left(2h + hPr - hNbPr - \right. \right. \\ & \left. \left. hNtPr - 6hPr\lambda - 3hPr\lambda 1 + 2hPr\lambda\lambda 1 + h(-2 + Pr\eta^2 + 4Pr\lambda + 2Pr\lambda 1 + \right. \right. \\ & \left. \left. 2\eta(-1 + 2Pr\lambda + Pr\lambda 1)) \right) \right), \dots \end{aligned} \quad (55)$$

$$\begin{aligned} \psi_1 = & \frac{1}{2Nb} (2Nbe^{-2\eta}(hLeNb(-1 - 2\lambda(-1 + \lambda 1) + \lambda 1) + e^\eta(2hNb + hLeNb \\ & + 2hNt - \\ & 6hLeNb\lambda - 3hLeNb\lambda 1 + 2hLeNb\lambda\lambda 1 + h(-2Nt(1 + \eta) + Nb(-2 + Le\eta^2 + \\ & 4Le\lambda + 2Le\lambda 1 + 2\eta(-1 + 2Le\lambda + Le\lambda 1))))), \dots \end{aligned} \quad (56)$$

$$\begin{aligned} \tau_1 = & \frac{1}{2}e^{-2\eta} \left(h(-2Pe + LbPr(-1 + 2\lambda + \lambda 1 - 2\lambda\lambda 1)) + e^\eta \left(-e^{-\eta}h(-2e^\eta - \right. \right. \\ & \left. \left. 2Pe - LbPr - 2e^\eta\eta + e^\eta LbPr\eta^2 + 2LbPr\lambda + 4e^\eta LbPr\lambda + 4e^\eta LbPr\eta\lambda + LbPr\lambda 1 + \right. \right. \\ & \left. \left. 2e^\eta LbPr\lambda 1 + 2e^\eta LbPr\eta\lambda 1 - 2LbPr\lambda\lambda 1 + 2e^\eta Pe\sigma + 2e^\eta Pe\eta\sigma) + h(LbPr\eta^2 + \right. \right. \\ & \left. \left. 2(-1 + LbPr(2\lambda + \lambda 1) + Pe\sigma) + 2\eta(-1 + LbPr(2\lambda + \lambda 1) + Pe\sigma)) \right) \right), \dots \end{aligned} \quad (57)$$

4 RESULTS AND DISCUSSION

The equations (9) - (12) with boundary conditions (13) are solved analytically by employing Homotopy Analysis Method and numerically by employing Runge-Kutta method. The obtained convergent solution depends on auxiliary linear operator L , auxiliary parameter h and the initial solution. For all variables h -curve is plotted to find the value of h and is used to get the convergent solution.

The significant parameters of this problem are Williamson parameter γ , Magnetic

field parameter M , Power exponent parameter λ , Brownian motion parameter Nb , Thermophoresis parameter Nt , Prandtl number Pr , Lewis number Le , Peclet number Pe , Bioconvection Lewis number Lb and Bioconvection parameter σ . The effect of these parameters on different profiles are as follows. In figure 2 we observed that velocity profile decreases with increase in γ . From figures 3 and 7 it is observed that increase in γ and Nt increases the temperature. From figures 4, 5, 6 and 8 we have observed that temperature profile decreases with increase in M, λ, Nb and Pr .

From figures 9, 10, 12 and 13 it is observed that the increase in values of λ, M, Nb and Le decreases the concentration profile, where as from figure 11 we can observe that increase in Nt increases the concentration profile. It is observed that increase in M, λ, Pr, Pe and Lb decreases the microorganism profile seen from figures 14, 15, 16, 17 and 18, where as increase in σ increases the microorganism profile shown in figure 19. Applied magnetic field will reduce temperature, concentration and gyrotactic microorganism in Williamson fluid.

We have obtained radius of convergence for velocity $R = 5.1342$, temperature $R = 7.0816$, concentration $R = 7.6923$ and gyrotactic microorganism $R = 7.6511$ by drawing Domb-Sykes plot in figures 20, 21, 22 and 23.

Comparison of numerical solutions and HAM solutions for velocity, temperature, concentration and gyrotactic microorganism profiles are shown from figures 24, 25, 26, 27 and in table 1 and observed good agreement. We compare our HAM solution with existing numerical solution of Haile [28] by the method Runge-Kutta integration technique. We have also solved numerically by R K Method and observe a very good matching in every case. We conclude that HAM gives almost exact solution.

5 Graphs and tables




Figure 1: Combine plot for h-curves




Figure 2: Effect of Williamson parameter γ on velocity profile




Figure 3: Effect of Williamson parameter γ on temperature profile




Figure 4: Effect of Magnetic field parameter M on temperature profile




Figure 5: Effect of power exponent parameter λ on temperature profile




Figure 6: Effect of Brownian motion parameter N_b on temperature profile




Figure 7: Effect of thermophoresis parameter N_t on temperature profile




Figure 8: Effect of Prandtl number Pr on temperature profile




Figure 9: Effect of power exponent parameter λ on concentration profile




Figure 10: Effect of Magnetic field parameter M on concentration profile




Figure 11: Effect of thermophoresis parameter Nt on concentration profile




Figure 12: Effect of Brownian motion parameter Nb on concentration profile




Figure 13: Effect of Lewis number Le on concentration profile




Figure 14: Effect of Magnetic field parameter M on gyrotactic microorganism profile




Figure 15: Effect of power exponent parameter λ on gyrotactic microorganism profile




Figure 16: Effect of Prandtl number Pr on gyrotactic microorganism profile




Figure 17: Effect of Peclet number Pe on gyrotactic microorganism profile




Figure 18: Effect of bioconvection Lewis number Lb on gyrotactic microorganism profile




Figure 19: Effect of bioconvection parameter σ on gyrotactic microorganism profile




Figure 20: Domb-Sykes Velocity Plot with $R=5.1342$




Figure 21: Domb-Sykes Temperature Plot with $R=7.0816$




Figure 22: Domb-Sykes Concentration Plot with $R=7.6923$




Figure 23: Domb-Sykes gyrotactic microorganism Plot with $R=7.0816$




Figure 24: Comparison of HAM solution and numerical solution for Williamson parameter $\gamma = 1$ on velocity profile




Figure 25: Comparison of HAM solution and numerical solution for Magnetic parameter $M = 0$ on temperature profile




Figure 26: Comparison of HAM solution and numerical solution for Lewis number $Le = 7$ on concentration profile




Figure 27: Comparison of HAM solution and numerical solution for Magnetic parameter $M = 0$ on gyrotactic microorganism profile

Table 1: Comparison of $f''(0)$ when $\gamma = M = 0$ and $h = 0.26156$

Parameter	$f''(0)$		
	Haile (R K integration technique) [28]	Present work (HAM)	Present work (R K Method)
λ_1			
0.1	0.46251	0.462509	0.462533
1	0.00000	0.0000	0.000000

REFERENCES

- [1] Pedley, T. J., Hill, N. A., and Kessler, J. O., The growth of bioconvection patterns in a uniform suspension of gyrotactic micro-organisms, *Journal of Fluid Mechanics*, vol.195, 1988, 223–237.
- [2] Pedley, T. J., Kessler, J. O., (1990). A new continuum model for suspensions of gyrotactic micro-organisms, *Journal of Fluid Mechanics*, vol.212, 1990, 155-182.
- [3] Pedley, T J, Kessler, J. O., Hydrodynamic Phenomena in Suspensions of Swimming Microorganisms, *Annual Review of Fluid Mechanics*, vol.24, 1992, 313-358.
- [4] Pedley, T. J., Instability of uniform micro-organism suspensions revisited, *Journal of Fluid Mechanics*, vol.647, 2010, 335–359.
- [5] Childress, S., Levandowsky, M. and Spiegel, E. A., Pattern formation in a suspension of swimming microorganisms, equations and stability theory. *Journal of Fluid Mechanics*, vol. 69(3), 1975, 591-613.
- [6] Talha, M. A., Osman Gani, M. and Ferdows, M., Numerical Solution of Steady Magnetohydrodynamic Boundary Layer Flow of Gyrotactic Microorganism for Williamson Nanofluid over Stretched Surface in the Presence of Exponential

- Internal Heat Generation, *International Journal of Mathematical and Computational Sciences*, vol.12, 2018, 39-47.
- [7] Alharbi, F. M., Naeem, M., Zubair, M., Jawad, M., Jan, W. U. and Jan, R., Bioconvection Due to Gyrotactic Microorganisms in Couple Stress Hybrid Nanofluid Laminar Mixed Convection Incompressible Flow with Magnetic Nanoparticles and Chemical Reaction as Carrier for Targeted Drug Delivery through Porous Stretching Sheet, *Molecules*, vol.26(13), 2021, 3954.
- [8] Bhatti, Muhammad M., Marin, Marin, Zeeshan, Ahmed, Ellahi, R. and Abdelsalam, Sara I., Swimming of Motile Gyrotactic Microorganisms and Nanoparticles in Blood Flow Through Anisotropically Tapered Arteries, *Frontiers in Physics*, vol.8, 2020, 1-9.
- [9] Lv, Y.-P., Algehyne, E. A., Alshehri, M. G., Alzahrani, E., Bilal, M., Khan, M. A., and Shuaib, M., Numerical approach towards gyrotactic microorganisms hybrid nanoliquid flow with the hall current and magnetic field over a spinning disk, *Scientific Reports*, vol.11(1), 2021.
- [10] Al-Khaled, Kamel., Khan, Sami Ullah., and Khan, Ilyas., Chemically reactive bioconvection flow of tangent hyperbolic nanoliquid with gyrotactic microorganisms and nonlinear thermal radiation, *Heliyon*, vol.6(1), 2020, 1-7.
- [11] Lund, Liaquat Ali., Omar, Zurni. and Khan, Ilyas., Analysis of dual solution for MHD flow of Williamson fluid with slippage, *Heliyon*, vol.5(3), 2019, 1-20.
- [12] Atif, S. M., Hussain, S., and Sagheer, M., Magnetohydrodynamic stratified bioconvective flow of micropolar nanofluid due to gyrotactic microorganisms, *AIP Advances*, vol.9(2), 2019, 025208.
- [13] Yusuf, T. A., Mabood, F., Prasannakumara, B. C. and Sarris, I. E., Magneto-Bioconvection Flow of Williamson Nanofluid over an Inclined Plate with Gyrotactic Microorganisms and Entropy Generation, *Fluids*, vol.6(3), 2021, 109.
- [14] Shahid, A., Zhou, Z., Bhatti, M. M., and Tripathi, D., Magnetohydrodynamics Nanofluid Flow Containing Gyrotactic Microorganisms Propagating Over a Stretching Surface by Successive Taylor Series Linearization Method, *Microgravity Science and Technology*, 2018.
- [15] Mahdy, A., Gyrotactic Microorganisms Mixed Convection Nanofluid Flow along an Isothermal Vertical Wedge in Porous Media, *International Journal of Aerospace and Mechanical Engineering*, vol.11, 2017.
- [16] Jayachandra Babu, M. and Sandeep, N., Effect of nonlinear thermal radiation on non-aligned bio-convective stagnation point flow of a magnetic-nanofluid over a stretching sheet, *Alexandria Engineering Journal*, vol. 55(3), 2016, 1931–1939
- [17] Khan, W.A., Makinde, O.D., Khan, Z.H., Non-aligned MHD stagnation point flow of variable viscosity nanofluids past a stretching sheet with radiative heat,

- International Journal of Heat and Mass Transfer, 2016, 525–534.
- [18] Chakraborty, Tanmoy., Das, Kalidas. and Kundu, Prabir Kumar., Framing the impact of external magnetic field on bioconvection of a nanofluid flow containing gyrotactic microorganisms with convective boundary conditions, Alexandria Engineering Journal, 2016.
- [19] Raju, C.S.K. and Sandeep, N., Heat and mass transfer in MHD non-Newtonian bio-convection flow over a rotating cone/plate with cross diffusion, Journal of Molecular Liquids, 2016, 115–126.
- [20] Bees, M. A. and Hill, N. A., Non-linear bioconvection in a deep suspension of gyrotactic swimming micro-organisms, Journal of Mathematical Biology, vol.38(2), 1999, 135–168.
- [21] Akbar, Noreen Sher., Bioconvection peristaltic flow in an asymmetric channel filled by nanofluid containing gyrotactic microorganism, International Journal of Numerical Methods for Heat and Fluid Flow, vol.25(2), 2015, 214–224.
- [22] Kuznetsov, A.V., The onset of nanofluid bioconvection in a suspension containing both nanoparticles and gyrotactic microorganisms, International Communications in Heat and Mass Transfer, vol. 37(10), 2010, 1421–1425.
- [23] Kuznetsov, A.V., Bio-thermal convection induced by two different species of microorganisms, International Communications in Heat and Mass Transfer, vol.38(5), 2011, 548–553.
- [24] Ahmad, S., Ashraf, M. and Ali, K., Nanofluid flow comprising gyrotactic microorganisms through a porous medium, Journal of Applied Fluid Mechanics, vol. 13, No. 5, 2020, 1539-1549.
- [25] Ahmad, S., Ashraf, M. and Ali, K., Heat and mass transfer flow of gyrotactic microorganisms and nanoparticles through a porous medium, International Journal of Heat and Technology, vol. 38, No. 2, 2020, 395-402.
- [26] Ramzan, Muhammad., Chung, Jae Dong. and Ullah, Naeem., Radiative magnetohydrodynamic nanofluid flow due to gyrotactic microorganisms with chemical reaction and non-linear thermal radiation, International Journal of Mechanical Sciences, 2017.
- [27] Acharya, Nilankush., Das, Kalidas. and Kundu, Prabir Kumar., Framing the effects of solar radiation on magneto-hydrodynamics bioconvection nanofluid flow in presence of gyrotactic microorganisms, Journal of Molecular Liquids, vol. 222, 2016, 28–37.
- [28] Haile, E. and Shnkar, B., Boundary-Layer Flow of Nanofluids over a Moving Surface in the Presence of Thermal Radiation, Viscous Dissipation and Chemical Reaction, Applications and Applied Mathematics, vol. 10, Issue 2, 2015, 952-969.
- [29] Liao, S.J., Homotopy analysis method in nonlinear differential equations, Springer, 2011.

- [30] Liao, S.J., *Advances in the homotopy analysis method*, World Scientific Publishing Co. Pte. Ltd., 2013.
- [31] Liao, S.J., *Beyond Perturbation: Introduction to the Homotopy Analysis Method*, Chapman and Hall/CRC Press, Boca Raton, 2003.
- [32] Liao, S.J., *On the homotopy analysis method for nonlinear problems*, *Journal of Applied Mathematics and Computation*, vol. 147, 2004, 499-513.
- [33] Liao, S.J., *Comparison between the homotopy analysis method and homotopy perturbation method*, *Applied Mathematics and Computations*, vol. 169, 2005, 1186-1194.
- [34] Liao, S.J., *Beyond perturbation, a review on the basic ideas of the homotopy analysis method and its applications*, *AdV Mech.*, 2008, 1-34.
- [35] Liao, S.J., *Notes on the homotopy analysis method: some definitions and theorems*, *Commun. Nonlinear Sci. Numer. Simulat.*, 2009, 983–997.
- [36] Rekha, K., Asha, C.S. and Achala, L.N., *Williamson Nanofluid Flow over a Moving Surface on Boundary Layer by Homotopy Analysis Method*, *International Journal of Mechanical Engineering*, vol.7, No.2, 2022, 1723–1736.
- [37] Achala, L.N. and Sathyanarayana, S.B., *Homotopy Analysis Method For Nonlinear Boundary Value Problems*, *JP of Applied Mathematics*, vol. 5, Issues 1 & 2, 2012, 27-46.
- [38] Achala, L.N., Madusudhan, R. and Sathyanarayana, S.B., *Homotopy Analysis Method To Solve Boussinesq Equations*, *Journal of Advances in Physics*, vol.10(3), 2015, 2825-2833.
- [39] Achala, L.N., Madusudhan, R. and Sathyanarayana, S.B., *Semi Analytic Approximate Solution Of Nonlinear Partial Differential Equations*, vol.4(6), 2016, 1418-1428.
- [40] Achala, L.N., Madusudhan, R. and Sathyanarayana, S.B., *Study Of Compressible Fluid Flow In Boundary Layer Region By Homotopy Analysis Method*, *International Journal of Latest Trends in Engineering and Technology*, vol.9(1), 2017, 028-039.
- [41] Sathyanarayana, S.B. and Achala, L.N., *Approximate Analytical Solution of Magnetohydrodynamics Compressible Boundary Layer flow with Pressure Gradient and suction/injection*, *Journal of Advances in Physics*, vol. 6(3), 2014.
- [42] Sathyanarayana, S.B. and Achala, L.N., *Approximate analytic Solution of Compressible Boundary Layer flow with an adverse Pressure Gradient by Homotopy Analysis Method*, *Theoretical Mathematics & Applications*, vol. 5(1), 2015, 15-31.

